

Report on the Biodiversity of area of Proposed Venkatesh Khani Coal Mine

of The Singareni Collieries Company Ltd. (SCCL) (A Government Company) 2022

CONTENTS

Background	1 - 4
Aim and Objectives	5
Study Area	6 - 14
Methodology	15 - 21
Results & Discussion	22 - 60
Conclusion	61
Recommendations	61
Photo plates	62 - 74
References	75

Summary

Initially the mining by underground method started in Kothagudem Area during 1937. In the early 70s, underground mines in the name of 8, 9, 10, 11 incline in an area of 408 Hectares were operational in the proposed study area, which were subsequently closed in 1992. From these underground mines, 5.56 million tons of coal have already been extracted. Subsequently, one opencast mine by the name of Goutham Khani (GKOC) started in the year 1993 in a total area of 902 Hectares and which has been closed during March 2022. This opencast was the conversion of the four underground mines, where the 74.29 million tons leftover of earlier underground operation. In total, 76.91 million tons (underground -5.56 + OC - 71.35) of coal has been removed from this area.

An average of manpower spent in these underground mines was around 6000 to 7000 and in the GK Opencast was of 1611 in total. Besides that, many coal loaded lorries/ vehicles move from mines to different places on daily basis (300 to 400 lorries per day).

Since, last 50 years of mine operations in this area, already movement of vehicles, people, constructions have started and as such hardly any area left undisturbed. The impact of the mining on the flora and fauna was also observed to be minimal, and records were also found that the fauna has moved away from this area to the nearby adjacent forest areas, plantation areas developed by SCCL and forest department in the 10 Kms radius from the proposed area. SCCL and Forest Department has taken many individual initiatives and joint actions in protecting, enhancing, and sustaining the biodiversity and its habitats in the adjoining forest areas in the buffer areas of mines to the 10 Kms radius. Hence the impact is very less.

Also, a National Highway (NH 30) connecting Kothagudem to Vijayawada passes through the forest and mine area, connected with many small public roads which is used by people actively. So, this also shows that there is not any fresh intrusion to the biodiversity/ wildlife by this proposed mine.

In this quarried area, SCCL has now proposed to dump the overburden material going to be excavated from the proposed VKOC to save fresh land.

Background

As demand for energy increases, in absence of sufficient alternative energy source, Coal mining will remain an important part of global, national, and local economies in the foreseeable future. To meet the ever-increasing demand for Coal and to achieve the targets fixed by the Government of India from time to time, SCCL has embarked upon large-scale expansion of coal mining to increase coal production. Venkatesh Khani Coal Mine is one of the new Project identified in Ramavaram RF, Ramavaram Range of Kothagudem Forest Division.

At present there are two Underground mines viz., Venkatesh Khani No.7 Incline (384.40 Ha. R.F) and PVK No.5 Incline (789.78 Ha. R.F) for which forest permission was obtained for underground rights for 1174.18 Ha. vide MoEF, Government of India vide Lr.No.8-277/85-FC, dated 17.02.2009 with validity for 20 years i.e., till 16.02.2029.

S. No.	Purpose of RF diverted	Extent (in Ha.)	MoEF&CC Letter No.	Validity till
1.	Venkatesh Khani No.7 Inc.	384.40	Lr.No.8-277/85-FC, dated 17.02.2009	16.02.2029
2.	Padmavathi Khani(PVK) No.5	789.78	-Do-	-Do-
3.	Surface use for installation of Borewell in existing Padmavathi Khani No.5 Inc.	2.85*	Stage-I clearance granted by MoEF&CC vide letter No.8- 277/1985 – FC (VOI), Dated: 22-06-2018.	Pending with the MoEF&CC, GoI/Forest dept., for issue of final approval.
	Total:	1174.18		*2.85 Ha. is the surface area of UG Mine of PVK-5 Incline.
4	Tella vagu (Stream) diversion to avoid seepage of water into UG Mine	11.96	MoEF&CC vide letter no. FCA/16.1/70/AP/MIN/07 dated.31.08.2005 Original course of stream was filled up with soil and planting done with local species as on today it is almost fully stocked. Further, there is no need to the re-divert/re-dig this stream. However, for the purpose of EC of total area of (1174.18+11.96) having mining lease. This is included in the total area of the Project.	

Similarly, permission under F (C) Act-1980, was also obtained for doing Opencast Mining on the name of Goutham Khani Opencast project (GKOCP), details of which are as follows:

S.No.	Purpose of RF diverted	Extent (in Ha.)	MoEF&CC Letter No.	Validity till
1.	1 st Renewal of GK OCP (Phase-I)	161.31	MoEF&CC Letter No.8-117/2002 – FC, Dated: 01-02-2010.	31.01.2030
2.	GK OCP (Phase- III)	154.96	MoEF&CC Letter No.8-62/2005 – FC, Dated: 09/15-07-2008.	14.07.2028
	Total:	316.27		

Similarly, an area of 124.965 Ha. was also diverted and it is under process with Forest department/ MoEF&CC, GoI, for renewal. Details of these diversions are as follows:

S.No.	Purpose of RF diverted	Extent (in Ha.)	MoEF&CC Letter No.
1.	1 st Renewal of GK OCP (Phase- II) (Extent increased as per DGPS Survey and Authentication by PCCF, Hyderabad).	124.065	MoEF&CC Letter No.8-17/98- FC, Dated: 08-02-1999 (for original diversion of 124.00 Ha.)
2.	Surface use for installation of Borewell Padmavathi Khani No.5 Inc.	0.90	1 st renewal of File No.8- 277/1985FC(vol. I) Dated: 15-10-2019(Stage-I Letter).
	Total:	124.965	

In the present proposal, it is proposed to utilize parts of the existing forest area of the UG Mines of VK-7 incline and PVK-5 Incline by converting parts of underground mining into opencast **and continuing part UG Mining in 469.6336 Ha. This amalgamated mine will be called Venkatesh Khani coal mine.** The details of area proposed for utilization in the new project from underground mines is as follows:

S.No.	Name of the UG Mine	Extent of RF land proposed for conversion (in Ha.)	RF Area proposed for continuation under UG	Total
1.	Venkatesh Khani No.7 Inc.	237.8569	146.5431	384.40
2.	Padmavathi Khani(PVK) No.5	411.4445	378.3355	789.78
		649.3014	524.8786	1174.18

The balance extractable coal reserves of PVK No.5 & VK No.7 Inc is 190.11 million Tons of which about 64.18 million Tons is in Forest Land in Compartment Nos.2 (part), 3(part), 4(part), 5(part) & 11(part) of Ramavaram RF, and the balance of 125.93 million Tons is in non-Forest land. However, mining (OC) will simultaneously start both in forest and non-Forest land as per the approved Mining Plan.

The proposed fresh diversion of forest land for surface use from underground to opencast is 649.3014 Ha for which this permission is requested. However, this area will be mainly used for quarrying and excavated material will be taken into the left-over quarry as well over burden dumps of the GKOCP for which already permission is available and at appropriate time (before two years after expiry of permission) proposal will be submitted for continuation of mining activities (dumping of excavated material of proposed Venkatesh Khani Coal Mine.

The total project area of **Venkatesh Khani Coal Mine** is 2403.17 Ha. This area is covered in 2 existing Mining Leases granted under MM (D&R) Act, 1957. The details of Mining Leases as per MMDR Act 1957 are as follows:

Fig in Ha.					
Name of Mining	Lease Area	NFL	FL		
Lease			Surface rights	UG rights	
Kothagudem Mining Lease (2 nd Renewal)	5158.00	3692.00	291.82	1174.18	
GK OC Mining lease(1 st renewal)	261.31	0.00	261.31	0.00	

The present proposed area is falling in Compartment Nos.2 (part), 3(part), 4(part), 5(part) & 11(part) of RF in Ramavaram R.F. of Kothagudem Division. The excavated material generated from the proposed OC mine will be transported through non-forest land for dumping into existing old mine of Goutham Khani Opencast. In the process the National Highway-30, passing through the non-forest land to the extent of 3.20 Kms will be diverted as per norms of the NHAI after obtaining suitable no objection certificate. Further, most of the forest area proposed for diversion either falls in to open forest, scrub forest or having grasses and degraded, wherever, better density is there it is because of plantations raised by the SCCL over the surface of the ongoing UG Mines.

Venkatesh Khani Coal Mine is planned to produce 6.3 (OC 5.3 + UG 1.0) million Tons of coal per annum (peak production) with a net capital out lay off about Rs.480.00 Crores and has a life of 40 years including pre-mining and post mining activities. The Mining plan including Mine closure plan has been approved by Ministry of Coal, Govt. of India vide Lr. No. 38011/12/2017-PCA, dated 27.01.2020. Further, the Nation will benefit by achieving increased coal production by exploiting coal reserves of the project and gets benefited financially by way of Royalty, excise Duty, Sales tax, and electricity supply continuously. Further, this project will also contribute benefit to the society by way of increased direct and indirect employment resources resultant improvement in standard of living and rise in per capital income of the population of surrounding villages of this area.

Generally, the activities of a mining could potentially pose some threats to biodiversity. A mining company must have a better understanding to appreciate the value of biodiversity in its long-term operations under its sustainable mining practice. It should consider all environmental impacts.

Habitat clearing during mining is generally unavoidable. However, impacts on fauna, including rare or threatened species, can be reduced through careful planning. Progressive clearing allows time for the animals to move into adjoining unmined/forest habitats. Despite the significant potential for negative impacts on biodiversity from mining operations, there are many opportunities for a mining company to enhance biodiversity conservation within their areas of operations. Being proactive in the assessment, monitoring, and management of biodiversity is important not only for new mine operations but also for those that have been operating for many years.

SCCL has been continuously adopting new and scientific approaches to managing biodiversity/wildlife as part of their commitments to establishing and maintaining the ecological balance. Prior to any mine expansion/operation in SCCL mines, the company reviews the biodiversity values and ensure that steps are taken to restore biodiversity wherever possible in scientific and sustainable manner. Removal of vegetation obviously has an immediate impact on local biodiversity, while ground disturbance may force biodiversity survival in rehabilitated areas.

Need of a baseline study:

A scientific intervention to assess the species diversity of an area which is selected for mining is necessary to understand the ecology, species composition and for enhancing the effectiveness of land management before and after mining.

Biodiversity is characteristically defined on three levels: genetic diversity, species diversity and ecosystem diversity. Biologists are aware of the importance of understanding diversity, at best with respect to the increasing loss of species due to the growing influence of human activities.

The term biodiversity describes the total number, variety, and variability of living organisms as well as the diversity of the ecosystem. Even though, there have been many different interpretations of diversity, the concept of biodiversity is the integration of biological variability across all scales, from genetic level, through species and ecosystems, to the landscapes that they form, or are part of, and the ecological processes that support them. The world's biodiversity includes all living organisms (animals, plants, fungi, and microbial groups inclusive of genetic diversity and ecosystem/landscape diversity) in their interactive state contributing to a multitude of services of relevance to sustain the ecological integrity for the benefit of the humankind. It defines biodiversity as the variability among living organisms from all sources including, inter alia (among other things), terrestrial and aquatic ecosystems and the ecological complexes of which they are a part; this includes diversity of any ecosystem is far too complex to be comprehensively quantified, suitable indicators or surrogates of biodiversity are needed.

Conceptually, species richness appears as the most intuitive and straightforward parameter to measure biodiversity. In addition, measures of biodiversity have been developed; the Shannon index (H'), and Simpson index (D) was considered measures used to compare diversity among communities. In general, most studies have been carried out on specific selected taxa only. Scientific biodiversity interventions will reveal data on availability of endemic/range-restricted species, whereby policies could be formulated for their future preservation. Such knowledge of the presence/absence of important species also facilitates potential conservation of critical microhabitats in and around the operational area.

Hence, as per the standard ToR condition "A detailed biodiversity assessment study of the proposed area" (core zone and buffer zone, 10 Kms radius of the periphery of the mine lease) has been carried out. Details of flora and fauna have been documented and furnished here in this report. This report carries information related to wildlife populations, habitat, and information gaps (baseline data) of the proposed mine areas and its adjoining forests areas.

Aim and Objectives

Aim:

There is an urgent need for interpreting this inventory before initiating mine operations and disseminating this information for sustainable management of this un-explored habitat and for encouraging further research and conservation in the adjoining forest areas, which will be the new home-range to the species getting disturbed/moving from this habitat.

Objectives:

For sustainable and eco-friendly mining project development, biological environment is studied. The objectives of the study are:

- 1. To register and validate existing information on biodiversity of the proposed area
- 2. To undertake field studies to study the impact of mines and record selected floral and faunal species including prominent invertebrates & soil surface arthropods
- 3. To recommend mitigation plans for species and habitat concern.
- 4. To assess any adverse impact on the local fauna because of the proposed mine.
- 5. Indicate the impacts and suggest suitable mitigation measures against the adverse impacts if any,

Study Area

State	: Telangana
District	: Bhadradri – Kothagudem
Forest Division	: Kothagudem
Forest Range	: Ramavaram
Forest Section (s)	: Ramavaram & Penagadapa
Forest Beat (s)	: Ramavaram, Penuballi and Gareebpeta (Ramavaram Section) Musalivarre West (Penagadapa Section)
Total study area	: 649.3014 Ha (Part -A = 334.7372 Ha + Part -B = 258.6898 Ha + Part -C = 55.8744 Ha)
Forest Type:	Tropical Dry Deciduous Forest – Mainly Coppice growth
Habitat Type	: Mixed-dry-deciduous forest type with wide variety of microhabitats, ranging from open grasslands, water bodies and hillocks, plantations, and woodlands to scrub lands.

It is proposed to open a new coal mining project under the name of Venkatesh Khani Coal Mine (VK Coal Mine) by amalgamating the three mines (*Gautham Khani OC*, *Venkatesh Khani -7 Incline* and Padmavathi Khani-5 Incline) with 6.3 (OC - 5.3& UG -1.0) MTPA capacity in mine lease area of 2403.17 ha (1568.22 ha Forest Land and 834.95 ha Non-Forest Land) located in Venkatesh Khani Village, Kothagudem town, District Bhadradri Kothagudem (Telangana State).

Physiographic Features of the study area:

To assess impacts of any set of activities, the environmental factors both within the core zone and buffer zone are to be analyzed. The core zone in this case covers project area while the buffer area covers the 10 Kms area around the core area.

Core zone: VK Coal Mine project area is of flat topography, Bolligutta hill stands out as a prominent landmark of this area along the northern boundary. The general topographic elevation of the project area varies from 160m above MSL to 100m above MSL. The south part of the project is covered by the existing GKOC project area.

Buffer zone: The buffer area is a plain terrain with relief of 160m in the south and 80m in the north and sloping towards north and northeast. A few hillocks located are Kanigiri gutta (432m) & an unnamed hillock (360m) in the northeast, Balusu gutta (328m) in the southeast, and Bolli gutta (340m) at the periphery of the project are existing in the buffer area.

Drainage:

Buffer zone: The buffer zone area drainage is drained by Murredu river. Edula vagu flowing in the western half and joins to Murredu river near Gollagudem village in the northern part of the buffer area. Tellavagu flows in the eastern part and joins Edulavagu near Penuballi village. The drainage of the area is dentritic to parallel type with a density of about 1.80 Kms/ sq.Kms. Singabhupalam Cheruvu a major irrigation tank is in the western part about 7 Kms away from the proposed VK Coal Mine project.

Core zone: There is no effective drainage developed in this area, due to sandy soil cover. Tellavagu, an ephemeral stream Tellavagu which is a tributary of Edula vagu is flowing in the northern part of the project area. Drainage is moderately developed, and a few ephemeral streams are flowing and join the Tellavagu. The drainage density of the block area is about 1.65 Kms/ sq.Kms.



Fig. 1. Google map showing VK Coal Mine along with study area



Fig. 2. Google map showing Part - A of the study area with Nature Trails (Padmavathi Khani Mine)



Fig. 3. Google map showing Part - B of the study area with Nature Trails (Venkatesh Khani 7 incline Mine)



Fig. 4. Google map showing Part - C of the study area with Nature Trails (Gautham Khani Mine)

Fig. 5. Google map showing Core and Buffer Zones of GK OC, VK-7 and PVK-5 in 10Kms of radius



The study area does not form a part of any National Park or Sanctuary or Critical Wildlife habitat. No protected area is situated in the vicinity or within the Impact area of 10 Kms radius. The proposed study area is an existing mine. No important wild animals are noticed in the area. There is no endangered Flora and Fauna within study area. In the Impact area the important Schedule-I fauna found are Varanus and Python.

List of Villages within the 10Kms of radius from the proposed study area

- 1. Sitarampuram
- 2. Anandapuram
- 3. Gurulagudem
- 4. Chinthalapadu
- 5. Patta Cheruvu
- 6. Gudipadu
- 7. Bangarujalu
- 8. Patha Palvancha
- 9. Palvancha
- 10. Sitaramesth Banjara
- 11. Chathakonda
- 12. Kothagudem
- 13. Penuballi
- 14. Gollagudem
- 15. Laxmidevipalli
- 16. Hemachandrapuram
- 17. Sarvaram
- 18. Chunchupalli
- 19. Ramavaram
- 20.Kotha Anjanapuram
- 21. Narasimha Nagar
- 22. Garibpeta
- 23. Garimellapadu
- 24. Mangapeta
- 25. Anjanapuram
- 26. Sujathanagar
- 27. Siripuram
- 28.Nimmalagudem
- 29. Sithampeta
- 30.Raghavapuram
- 31. Thungaram
- 32. Repallevada
- 33. Arlapadu
- 34.Ganugapadu
- 35. Ravikampadu
- 36. Chandraguda
- 37. Ayyannapalem
- 38.Thippanapalli
- 39. Sitayegudem
- 40.Mohammad Nagar
- 41. Satyanarayanapuram
- 42.Rampuram
- 43. Penagadapa
- 44. Rudrampur
- 45. Laxmipuram Thanda
- 46.Dhanbad
- 47.4 Incline
- 48.Mulkagudem

pg. 12

- 49.Chaparalapalli
- 50.Kommagudem

Fig. 6. Map showing the forest compartments in the Core and Buffer Zones of GK OC, VK-7 and PVK-5 in 10Kms of radius





Fig. 7. Map showing the Canopy Density in the study area (Part A, B&C)

Methodology

The study was conducted in all the three proposed parts (A, B & C) (Fig. 2, 3 & 4) for mining lease. The varying habitats in these three parts give rise to diverse ecosystems providing nesting and roosting site for birds, mammals, reptiles, amphibians, insects, and butterflies.

The present study is an assessment of biodiversity in the proposed area over a period of about one year and again re-visited the area during October- November 2022.

The procedure therefore involved field surveys of selected components of biodiversity such as plants butterflies, amphibians, reptiles, birds, and mammals. Intensive survey methods (including night trails in the proposed area) were used involving Forest Beat Officers. The team walked the pre-identified nature trails (Fig. 2, 3 & 4) in the study area extensively covering the entire proposed study area to enumerate different species as and when they were encountered or sighted.

Studies on faunal diversity were performed during both diurnal and nocturnal timings. Point counts on the nature trials were performed to document the bird species, Opportunistic surveys and listing encounters in random trails were also performed.

All observations were conducted on foot and were aided by 80x40 binoculars. Photo records were obtained with the help of Nikon 24-3000 mm camera with 125x wide optical zoom lens.

Species were identified using

- "A pictorial guide to the mammals of the India". Avifauna species were identified using "A pictorial guideto the birds of the Indian Sub-Continent".
- Reptiles and amphibians were identified with the available identification keys include Fauna of British India.
- Filed guides and published literature were used for theidentification of Odonates, butterflies, arachnids, fishes, and other invertebrates.

Meteorological data:

Basic field conditions were recorded by observing visually. The collected data was compared with the online data portals providing meteorological data.

Floral sampling procedures:

Secondary data obtained from the Environment Management Plan already existing with the company was obtained and presented here in the report. (Fig. 7).

Faunal sampling procedures:

1. Pre-existing tracts and roads were marked as Nature Trails in each Part of the study area covering microhabitats within it (Fig. 2, 3 & 4). These tracts were walked during morning and evening hours and sampling was done using line transect method.

A total of 11 nature trails with 1100 mts each were walked during the survey. (Table.1). Six trails in Part – A, four trails in Part – B, and one Trail in Part – C, covering all the vegetation and habitat type in the proposed study area. (Fig. 2, 3 & 4).

	STAR	TING	END	DISTANCE	
TRAILS	LAT	LONG	LAT	LONG	IN METERS
1.	N17°30'14.598"	E80°39'29.022"	N17°30'38.128"	E80°39'51.403"	1100
2.	N17°28'59.555"	E80°40'20.490"	N17º29'28.267"	E80°40'40.853"	1100
3.	N17º28'58.613"	E80°40'20.907"	N17º29'12.146"	E80°40'55.242"	1100
4.	N17º28'39.982"	E80°39'38.146"	N17º28'23.562"	E80°39'41.339"	1100
5.	N17º29'38.4"	E80°39'32.526"	N17°30'11.703"	E80°39'39.723"	1100
6.	N17°30'10.379"	E80°40'22.720"	N17 ⁰ 30'31.013"	E80°40'31.786"	1100
7.	N17°29'41.580"	E80°40'25.348"	N17°29'41.328"	E80°40'48.795"	1100
8.	N17º30'8.180"	E80°40'18.141"	N17º30'34.688"	E80°40'0.076"	1100
9.	N17°29'29.789"	E80°39'37.310"	N17°29'47.372"	E80°40'4.776"	1100
10.	N17°29'55.422"	E80°40'7.0173"	N17º29'55.960"	E80°39'37.669"	1100
11.	N17º28'37.455"	E80°40'17.723"	N17º28'53.052"	E80°40'48.081"	1100

Table 1. Geographic coordinates nature trails in the PART A, B & C study area

Line Transect Method

The line transect method of Gaston (1975) is adopted in most of the population estimation studies.

Procedure: At the start of the transect, the transect name, transect bearing, locality, date, starting time and other details were noted in proforma, especially designed for the purpose. Then the team walked along the transect carefully and silently looking for animals on both sides of the transect. On sighting an animal or group of animals, the number is counted, and their position is noted. Then, their angular distance i.e., the distance from the observer to the animal was measured visually. The sighting angle was derived from the compass bearing of the animal's location and the transect compass bearing. The perpendicular distance of the animal from the transect was estimated by multiplying the angular distance with Sin of the sighting angle. The data obtained from different nature trails were pooled and mean perpendicular distance was estimated.

Then, the density of the animal was calculated using the following formula,

Density = n/2l*r

Where, n = number of animals sighted l = total length of the transect r = Angular distance

- 2. Mammalian herbivore presence was assessed through direct sighting only along the trails, and around watering points.
- 3. Carnivore presences were also studied through direct sighting only.
- 4. **Direct sighting and vocalizations** were recorded to study bird presence and validate existing data on birds of the all the Parts in the study area. Early morning and evening studies were carried out when bird activity is at its peak. The point-count method was used on the nature trails by marking each point at 200mts interval along each trail and the bird count was performed within a radius of 25mts at each point. (Fig. 8.). A total of 5 points were marked and used for the bird point-count at each trail in all the three parts of the study area (Table. 2). We collected

the data from 55 points established systematically and have been marked in all 11 nature trails of 1100mts each in the study area. (Table 2)





Bird Point Count Method

Procedure: A point count consists of standing in a specific location and counting birds. One counted the number of individual birds (of each species) within a circle of 20mts radius. When gathering data to compare one point count to the next, radius size was kept consistent. The radius of 20mts was as large as possible kept in the study area based on the vegetation type (mostly open forest) to maximize information gathering, and the birds could not be seen are heard and recorded throughout the nature trails in the study area. The 20mts radius point count at each point was performed to record all the birds seen or heard within a 5-minute period. We have recorded all birds that are seen or heard within the normal 20mts radius. At the same time, we have also recorded the birds that are heard or seen within a selected distance outside this 20mts radius. The birds counted within the 20mts radius were marked in the "Number of Birds" column of the data sheet and the birds seen or heard outside the 20mts radius (within a specified distance) are marked in the "Outside" column of the data sheet. Counting the same bird twice during the survey was avoided.

A bird that moves from the 0 - 20mts area was not counted twice. Counted the birds where it first appeared or sighted. The purpose of surveying birds in this manner is that it allows one to compare across sites.

Fig. 9. Diagrammatic representation of bird point-count design



Table 2. Geographic coordinates of bird point-count points marked in the naturetrails in all the PARTs A, B & C of the study area

Point No.	oint No. Latitude			Trail No.			
	Deg.	Min.	Sec.	Deg.	Min.	Sec.	
1	17	30	18.85598	80	39	34.08737	TRAIL-1
2	17	30	24.67052	80	39	36.66052	
3	17	30	30.91033	80	39	38.53268	
4	17	30	33.07393	80	39	44.58386	
5	17	30	35.00443	80	39	50.55766	
6	17	29	5.08251	80	40	23.53231	TRAIL-2
7	17	29	10.25375	80	40	27.62735	
8	17	29	15.67464	80	40	31.34328	
9	17	29	21.37146	80	40	34.45858	
10	17	29	26.24166	80	40	38.56585	
11	17	29	1.19389	80	40	27.01192	TRAIL-3
12	17	29	3.57506	80	40	33.31775	
13	17	29	6.04545	80	40	39.59193	
14	17	29	8.59187	80	40	45.83286	
15	17	29	11.1573	80	40	52.06552	
16	17	28	34.17069	80	39	41.11204	TRAIL-4
17	17	28	28.39292	80	39	44.23322	
18	17	28	22.61514	80	39	47.35436	
19	17	28	18.32158	80	39	47.712	
20	17	28	21.37383	80	39	43.75726	
21	17	29	44.00492	80	39	32.96772	TRAIL-5
22	17	29	50.47755	80	39	32.36692	
23	17	29	56.85014	80	39	33.10816	
24	17	30	3.02317	80	39	35.25616	
25	17	30	8.97316	80	39	37.93188	
26	17	30	11.83627	80	40	29.0699	TRAIL-6
27	17	30	16.12002	80	40	34.13094	
28	17	30	19.33687	80	40	39.49441	
29	17	30	23.89631	80	40	38.67032	
30	17	30	28.66663	80	40	34.05654	
31	17	29	42.04538	80	40	31.44358	TRAIL-7
32	17	29	43.25359	80	40	37.26235	
33	17	29	42.75506	80	40	43.93841	
34	17	29	42.3786	80	40	49.78611	
35	17	29	47.57437	80	40	53.87023	
36	17	30	12.89298	80	40	14.26144	TRAIL-8

37	17	30	18.38437	80	40	12.59121	
38	17	30	23.12239	80	40	8.92722	
39	17	30	26.85979	80	40	3.71859	
40	17	30	33.02159	80	40	2.18556	
41	17	29	34.25556	80	39	42.03252	TRAIL-9
42	17	29	36.4816	80	39	47.27233	
43	17	29	37.6769	80	39	53.8697	
44	17	29	40.31582	80	40	0.05682	
45	17	29	44.22869	80	40	3.9016	
46	17	29	56.47055	80	40	0.40638	TRAIL-10
47	17	29	58.69132	80	39	54.07332	
48	17	29	59.93205	80	39	47.43811	
49	17	30	1.8855	80	39	41.05976	
50	17	29	59.19858	80	39	37.43828	
51	17	28	41.15651	80	40	22.38711	TRAIL-11
52	17	28	42.92922	80	40	28.6734	
53	17	28	45.98556	80	40	34.26492	
54	17	28	48.81467	80	40	40.07795	
55	17	28	50.98344	80	40	45.74362	

The following details provide the protocol adopted for bird count:

- a. Selection of points: The study are satellite images were captured, and the bird points at 200mts distance each was fixed on the image having the pre-existing nature trails covering the vegetation sampling grids. The centre point of each point has been chosen based using GPS and systematic sampling was followed.
- b. Duration and time of counts: Counts were done within three hours after sunrise. This is when birds are most active. Also done night-time surveys to count nocturnal species (done within three hours after sunset). During point counts, recorded all birds seen and heard within the survey area. The team initially stood for two minutes to allow some settling time for the birds that were disturbed. Observations were recorded for five minutes.
- c. Recording method: For each sighting, the number of individuals, and the sighting distance was recorded. Flying birds and calls of the birds were recorded separately.
- *d.* Counting FO (Flyovers): All higher-flying birds (above the tallest structure in the study area) were also noted if they are within the boundaries of the point count area.
- e. Counting Birds Outside of Survey Area: Only birds seen or heard within the point count area were recorded.
- f. Estimating Abundance: When multiple sightings of a species occurred within a point count, we have included only multiple entries for a species of sure that they are different individuals.
- *g.* Avoided Artificial Densities: No sounds used that can attract birds to our trail or in the study are. No recorded calls, or any other methods that encourage birds to show themselves was used.

- *h.* **Set-up:** *Before conducting the point count, the boundaries of the area marked visually by using some identifiable object (e.g., a large tree). Also, marked the centre spot where we stood and observed the birds with GPS. The exact place was used each time for the survey.*
- *i.* Distance estimation: Ocular estimates of the distance were taken.
- *j.* Weather: While conducting the point count, the general climatic conditions were recorded; Wind intensity (calm, slight, gusty, strong) and sky condition (clear, cloudy). This is important because climatic variables are known to affect bird activity. Bird point count studies were avoided on rainy days or extremely windy days because birds don't produce calls during that time, hence affecting the detectability.
- 5. **Reptiles and amphibians** were recorded in through incidence rate along nature trails as well as occurrence and interception rates along water holes. Other than the day counts, sampling was also conducted during the early evening and night, when incidence rates are expected to be highest.
- 6. **Invertebrates** were estimated through incidence rates and direct sighting. Butterflies, ants, spiders, dragonflies, scorpions, bees, wasps, beetles, grasshoppers, and other prominent invertebrates were check-listed. Where identification is not possible in the field, photographs were taken for taxonomical purpose.
- 7. **Special habitats and eco-tones**, especially fields and grasslands were surveyed for estimating the presence of grassland birds, rodents, and small mammals.

Source of secondary information on wildlife:

- Secondary information on flora, fauna and their distribution were collected from Forest Department, Kothagudem (Working Plan/Management Plan of Kothagudem) Division, 2014-2024 of Bhadradri Kothagudem district. Also, data were collected on wildlife census conducted, incidents of forest fires, man-animal conflicts, compensation cases/details, water sources (natural and man-made structures), density enrichment plantations by Forest Department and SCCL, grassplots developed.
- Faunal availability was also verified from the local people and staff working in the forest department.
- Conservation status of the flora and fauna were cross checked under IUCN Red list through published literature and online as well as Indian Wildlife Protection Act (IWPA), 1972 and further Amendments.

Source of secondary information on villages and livestock:

The methodology adopted for the study mainly includes review of published secondary data and primary field survey (District Census Statistical Handbooks- 2011 and Primary Census Abstract of Census-2011) with respect to population, sex ratio, average household size, density of population, social stratification, literacy rate and occupational structure for 10 Kms radius study area. The villages falling within the study area were visited to have first-hand information on the socioeconomic conditions of the people living in these villages.

Traffic Study:

The traffic study has been conducted to know the prevailing traffic volumes on the existing approach roads to the project area. Even though no coal transport is proposed through road mode in the project. It is essential to consider these details for assessing the anticipated future traffic volumes as a part of overall impacts assessment for the project.

Methodology adopted for studying the Socio-economic Status within the 10Kms radius from the proposed study area:

The methodology adopted for the study mainly includes review of published secondary data and primary field survey (District Census Statistical Handbooks- 2011 and Primary Census Abstract of Census-2011) with respect to population, sex ratio, average household size, density of population, social stratification, literacy rate and occupational structure for 10 Kms radius study area. The villages falling within the study area were visited to have firsthand information on the socioeconomic conditions of the people living in these villages.

The information gathered during the interview and Participatory Rural Appraisal (PRA)/ Rapid Rural Appraisal (RRA) covered different aspects of socio-economic indicators to evaluate the present quality of life of the people living in the area and to know the people's perception on the proposed project.

Results & Discussion

Meteorological Data:

Kothagudem area experiences typical tropical climate of a distinct hot summer from March to May with occasional dust storms, a good monsoon between middle of June and September and a pleasant winter from December to February. The temperature varies between 9.10 C and 48.60 C.

Mon	Tue	Wed	Thu	Fri	Sat	Sun
			1	2	3	4
			+86" right+81"	+90°	+91°	+91°
5	6	7	8	9	10	11
+91°	+91°	+86°	+82"	+82° nigni+79*	+82°	+82°
12	13	14	15	16	17	18
+86°	+88°	+88°	+88°	+90°	+90° night+79*	+84°
19	20	21	22	23	24	25
84°	+84° night+79	+88°	+86° nigrit+79*	+84° night+79°	*86°	+88° right+79*
26	27	28	29	30		
+86°	+79° night+79*			+84" right+79*		
light+79*	night+79*			night+79*		

Fig. 10. Average Weather Data of the Study Duration (September 2022)



Fig. 11. Temperature Graph of the Study Duration

https://world-weather.info/forecast/india/kothagudem_7/september-2022/ https://www.accuweather.com/en/in/kothagudem/186810/septemberweather/186810

Three seasons studies were carried out at the time of preparation of EMP. Again, study was carried out during October- November 2022. The season during the study period was post-monsoon season in the study area. Most of the days were sunny and had rains in the evening. The wind was also light during the study period. Sky was recorded always clear and cloudy during the rains. The weather conditions were favorable during the study period to record the existing species in the proposed study area.

Floral Diversity:

A total of 313 plant species belonging to 241 genera and 83 families were documented in the study area (Core and Buffer zone). Among 313 species listed: 89 species are trees, 41 shrubs, 14 lianas, 22 climbers, 117 herbs, 29 aquatic herbs and one is epiphyte.

As the vegetation analysis suggests there are more herbaceous species (117 species) in comparison to tree species or other life forms as shown in figure.

This infers to the study area of proposed project is medium density forests.

Among all 313 species, 23 species are recorded in core zone, 122 species in buffer zone and 168 are recorded in both core and buffer zone areas of the study area. The species listed in core zone includes 5 trees, 3 shrubs, 1 climber and 14 herbs; in buffer zone 19 trees, 14 shrubs, 4 lianas, 11 climbers, 48 herbs and 26 aquatic herbs; in core and buffer 65 trees,24 shrubs, 10 lianas, 10 climbers, 55 herbs, 3 aquatic herbs and one is epiphyte.

S. No.	Scientific Name	Family	Habit	Vernacular Name	Study Zone	IUCN Status
1	Abildgaardia ovata	Cyperaceae	Н	-	B	LC
2	Abrus precatorius	Fabaceae	C	Guruginia	C&B	NA
3	Abutilon hirtum	Malvaceae	S	Palabenda	C&B	NA
4	Abutilon indicum	Malvaceae	S	Thuttutubenda	B	NA
5	Acalypha capitata	Euphorbiaceae	S	Chiru Kuppinta	C	NA
6	Acalypha indica	Euphorbiaceae	H	Muripinda	C&B	NA
7	Achyranthes aspera	Amaranthaceae	Н	Uttareni	B	NA
8	Acmella paniculata	Asteraceae	H	Maratimoggalu	C&B	LC
9	Actinoscirpus grossus	Cyperaceae	AH	-	B	NA
10	Adina cordifolia	Rubiaceae	Т	Rudraganapa, Patchabotruka	C&B	NA
11	Aegle marmelos	Rutaceae	Т	Maredu, Bilvumu	C&B	NA
12	Afrohybanthus enneaspermus	Violaceae	Н	Ratna purusha	С	NA
13	Agave americana	Asparagaceae	S	Kathaichettu	В	NA
14	Ageratum conyzoides	Asteraceae	Η	Adavipudeena	C&B	NA
15	Ailanthus excelsa	Simaroubaceae	Т	Pedda manu	C&B	NA
16	Alangium salviifolium	Cornaceae	Т	Nallaoodaga	C&B	LC
17	Albizia amara	Leguminosae	Т	Narlingi, Chigara	C&B	LC
18	Albizia odoratissima	Leguminosae	Т	Chinduga	C&B	LC
19	Albizia thompsonii	Leguminosae	Т	-	C&B	NT
20	Alternanthera pungens	Amaranthaceae	Н	Mulla Ponnaganti	C&B	NA
21	Alternanthera sessilis	Amaranthaceae	Η	Adavi Ponnaganti	C&B	LC
22	Alysicarpus hamosus	Leguminosae	Н	-	С	NA
23	Alysicarpus monilifer	Leguminosae	Н	Poosala mokka	В	NA
24	Amaranthus spinosus	Amaranthaceae	Η	Mundla	C&B	NA
25	Amaranthus viridis	Amaranthaceae	Н	Chilaka thotakura	В	NA
26	Ammannia baccifera	Lythraceae	AH	Nela Citramulam, Agnivendramu	В	LC
27	Ammannia multiflora	Lythraceae	AH	-	В	LC
28	Andrographis echioides	Acanthaceae	H	-	C&B	NA
29	Andrographis paniculata	Acanthaceae	H	Belavemu	C&B	NA
30	Annona squamosa	Annonaceae	Т	Seetaphalam	C&B	LC
31	Aponogeton crispus	Aponogetonaceae	AH	Nammapuvvu	В	LC
32	Aristida adscensionis	Poaceae	H	Cheepuru	C&B	NA
33	Aristida hystrix	Poaceae	Н	Chinnameesala gaddi	C&B	NA
34	Aristida setacea	Poaceae	Н	Chipurugaddi	C&B	NA
35	Aristolochia indica	Aristolochiaceae	C	Iswaraveru	C&B	NA
36	Asparagus racemosus	Asparagaceae	C	Sathavari	C&B	NA
37	Ayenia herbacea	Malvaceae	H	Magasirigadda	C&B	NA
38	Azadirachta indica	Meliaceae	Т	Vepa chettu	C&B	LC

Table 3. List of species of floral species documented

39	Bacopa monnieri	Plantaginaceae	AH	Brahmmi	В	LC
40	Balanites aegyptiaca	Zygophyllaceae	Т	Ingudi,	С	LC
				Garachettu		
41	Bambusa bambos	Poaceae	T	Veduru	C&B	NA
42	Barleria prionitis	Acanthaceae	H	Mullagorinta	B	NA
43	Bauhinia racemosa	Leguminosae	Т	Arichettu	C&B	NA
44	Bergia capensis	Elatinaceae	AH	Neerupavila	B	
45	Blepharis integrifolia	Acanthaceae	H			NA
46	Blepharis maderaspatensis	Acanthaceae	H	Anthritapoolu	C&B	NA
47	Blumea axillaris	Compositae	H	Киккародаки		NA
48	Biumea iacera	Nustaginagaaa	Н	- Dura oraș	D D	NA NA
49	Boernavia diffusa	Nyctaginaceae	Н	Atikimamidi	В	NA
50	Boerhavia erecta	Nyctaginaceae	Н	Pandari punarnava	В	NA
51	Bombax ceiba	Malvaceae	Т	Buruga	В	LC
52	Bonnaya ciliata	Linderniaceae	AH	-	В	LC
53	Boswellia serrata	Burseraceae	Т	Anduga	В	NA
54	Brachypterum scandens	Leguminosae	L	Nallateega	C&B	NA
55	Bridelia montana	Phyllanthaceae	S	Adavi jama	C&B	NA
56	Bridelia retusa	Phyllanthaceae	Т	Mullumaddi	C&B	LC
57	Buchanania cochinchinensis	Anacardiaceae	Т	Chinna morli	C&B	NA
58	Butea monosperma	Leguminosae	Т	Moduga	C&B	LC
59	Butea superba	Leguminosae	L	Teega modiga	C&B	NA
60	Cajanus scarabaeoides	Leguminosae	C	-	C&B	LC
61	Calotropis gigantea	Apocynaceae	S	Tella Jilledu	В	NA
62	Calotropis procera	Apocynaceae	S	Erra jilledu	В	NA
63	Canavalia gladiata	Leguminosae	L	Thammakaya	В	NA
64	Canscora heteroclita	Gentianaceae	AH	Thambakaya	В	NA
65	Canthium coromandelicum	Rubiaceae	S	Sinnabalusu, Balusu	C&B	NA
66	Capparis sepiaria	Capparaceae	S	Nallauppi	В	LC
67	Capparis spinosa	Capparaceae	L	-	C&B	LC
68	Careya arborea	Lecythidaceae	Т	Budadarmi	C&B	NA
69	Carissa carandas	Apocynaceae	S	Kalivi	В	NA
70	Cassia fistula	Leguminosae	Т	Rela chettu	C&B	LC
71	Cassytha filiformis	Lauraceae	C	Pasuputivva	C&B	NA
72	Catunaregam spinosa	Rubiaceae	S	Manga	C&B	NA
73	Celastrus paniculatus	Celastraceae	L	Jyothismathi, Teegapalleru	C&B	NA
74	Ceratopteris thalictroides	Pteridaceae	AH	-	В	LC
75	Chara globularis	Characeae	AH	Chara	В	LC
76	Chloris barbata	Poaceae	Н	Jada kunchula gaddi	C&B	NA
77	Chloroxylon swietenia	Rutaceae	Т	Billudu	C&B	VU
78	Chromolaena odorata	Compositae	S	Porangi	C&B	NA
79	Chrozophora rottleri	Euphorbiaceae	Н	Lingamirapa	В	NA
80	Chrysopogon fulvus	Poaceae	Н	-	С	NA
81	Cleistanthus collinus	Phyllanthaceae	Т	Kodise	C&B	VU
82	Cleome viscosa	Cleomaceae	Н	Kukka vominta	C&B	NA
83	Clitoria ternatea	Leguminosae	С	Dintena teega	В	NA
84	Coccinia grandis	Cucurbitaceae	C	Donda	В	NA

85	Cocculus hirsutus	Menispermaceae	C	C Chinnadusar C&B		NA
86	Coldenia procumbens	Boraginaceae	Н	Bukkinaaku	C&B	LC
87	Colocasia esculenta	Araceae	Н	Chemadumpa	В	LC
88	Combretum albidum	Combretaceae	L	Yadateega	В	NA
89	Commelina benghalensis	Commelinaceae	AH	Vennadevikura	C&B	LC
90	Commelina imberbis	Commelinaceae	Н	-	C&B	LC
91	Cordia dichotoma	Boraginaceae	Т	Iriki	В	LC
92	Crateva adansonii	Capparaceae	Т	Uskimanu,	В	LC
93	Crotalaria hebecarpa	Fabaceae	Н	-	C&B	NA
94	Croton bonplandianus	Euphorbiaceae	Н	Vanamokka	В	NA
95	Cryptolepis buchananii	Apocynaceae	С	Adavipalateega	С	NA
96	Curculigo orchioides	Hypoxidaceae	Н	Nelathadi	C&B	NA
97	Curcuma pseudomontana	Zingiberaceae	Н	Adavipasupu	C&B	VU
98	Cyanotis axillaris	Commelinaceae	AH	Neelavanthi	C&B	LC
99	Cyanthillium albicans	Compositae	Н	Garitakamma	C&B	NA
100	Cyanthillium cinereum	Compositae	Н	Sahadevi	В	NA
101	Cycas rumphii	Cycadaceae	Т	Ranhaguvva	С	VU
102	Cymbopogon coloratus	Poaceae	Н	Bodagaddi	C&B	NA
103	Cymbopogon martini	Poaceae	Н	Kamaanchi kasuvu	C&B	NA
104	Cynodon dactylon	Poaceae	Н	Garika	C&B	NA
105	Cyperus corymbosus	Cyperaceae	AH	-	В	NA
106	Cyperus difformis	Cyperaceae	AH	-	В	LC
107	Cyperus exaltatus	Cyperaceae	AH	-	В	LC
108	Cyperus michelianus	Cyperaceae	AH	-	В	LC
109	Cyperus mindorensis	Cyperaceae	AH	Gandala	C&B	NA
110	Cyperus pangorei	Cyperaceae	AH	-	В	LC
111	Cyperus rotundus	Cyperaceae	Н	Thunga	В	LC
112	Dalbergia lanceolaria	Leguminosae	Т	Patchari	C&B	NA
113	Dendrocalamus strictus	Poaceae	Т	Sadanam	C&B	NA
114	Dendrophthoe falcata	Loranthaceae	S	Jiddu, Yolinga	C&B	NA
115	Dentella repens	Rubiaceae	Н	Katakura	В	LC
116	Dentella repens var. serpyllifolia	Rubiaceae	Н	-	В	NA
117	Dichanthium annulatum	Poaceae	Н	Errasangali gaddi	В	NA
118	Dichrostachys cinerea	Leguminosae	S	Velthuruchettu	C&B	LC
119	Dioscorea pentaphylla	Dioscoreaceae	C	Adaviginasu teega	C&B	NA
120	Diospyros chloroxylon	Ebenaceae	Т	Ullinda	C&B	NA
121	Diospyros melanoxylon	Ebenaceae	Т	Beediakulu, C&B Thuniki		NA
122	Diospyros montana	Ebenaceae	Т	Muchha thuniki	В	NA
123	Dodonaea viscosa	Sapindaceae	S	Bandaru, Pullivavili	C&B	LC
124	Drimia indica	Asparagaceae	H	Adavi ulli	С	NA
125	Drypetes sepiaria	Putranjivaceae	S	Putrajivika, Kuduru	В	NA
126	Eclipta prostrata	Asteraceae	Н	Guntagalagara	C&B	LC
127	Ehretia aspera	Boraginaceae	Т	Paldattam	C&B	DD
128	Eleocharis geniculata	Cyperaceae	AH	-	В	LC

129	Elytraria acaulis	Acanthaceae	H	Nela marri B		NA
130	Eragrostiella bifaria	Poaceae	Н	Noolugaddi	В	NA
131	Eragrostis tenella	Poaceae	Н	Chinna garikagaddi	C&B	NA
132	Eragrostis viscosa	Poaceae	Н	-	C&B	NA
133	Eriocaulon quinquangulare	Eriocaulaceae	Н	-	С	NA
134	Erythroxylum monogynum	Erythroxylaceae	S	Gatiri, Adavi gongura	C&B	NA
135	Euphorbia hirta	Euphorbiaceae	H	Nanubalu	C&B	NA
136	Euphorbia nivulia	Euphorbiaceae	Т	Aakujemudu, Bonthajemudu	В	NA
137	Euphorbia thymifolia	Euphorbiaceae	Н	Reddivaari nanambralu	В	NA
138	Evolvulus alsinoides	Convolvulaceae	H	Vishnukrantha	В	NA
139	Evolvulus nummularius	Convolvulaceae	Н	-	C&B	NA
140	Ficus benghalensis	Moraceae	Т	Marri	C&B	NA
141	Ficus hispida	Moraceae	S	Bemmedu akulu	C&B	LC
142	Ficus mollis	Moraceae	Т	Konda kalijuvvi	C&B	NA
143	Ficus racemosa	Moraceae	Т	Medi	C&B	LC
144	Ficus religiosa	Moraceae	Т	Ravi chettu	В	NA
145	Fimbristylis aestivalis	Cyperaceae	AH	-	В	NA
146	Fimbristylis argentea	Cyperaceae	AH	-	В	LC
147	Fimbristylis quinquangularis	Cyperaceae	AH	-	В	LC
148	Flacourtia indica	Salicaceae	S	Nakka neredu	C&B	LC
149	Galactia striata	Leguminosae	C	-	В	NA
150	Gardenia gummifera	Rubiaceae	Т	Bikki	C&B	LC
151	Gardenia latifolia	Rubiaceae	Т	Peddabikki	C&B	NA
152	Garuga pinnata	Burseraceae	Т	Garuga, Kondavepa	В	NA
153	Getonia floribunda	Combretaceae	L	Pootangiteega	C&B	NA
154	Gisekia pharnaceoides	Gisekiaceae	H	Isukadantikura	В	NA
155	Givotia moluccana	Euphorbiaceae	Т	Tella Poliki C&B		NA
156	Glinus lotoides	Molluginaceae	Н	Tellaporaku	В	LC
157	Glinus oppositifolius	Molluginaceae	Н	Chayunta rashiaku	В	LC
158	Gloriosa superba	Colchicaceae	Н	Naabhi, Adavinaabhi	С	LC
159	Gmelina arborea	Lamiaceae	Т	Gummaditeku, Gambharee	C	LC
160	Gmelina asiatica	Lamiaceae	L	Chinna adavigummadi	Chinna B adavigummadi	
161	Gomphrena serrata	Amaranthaceae	Η	Bendumalli	В	NA
162	Grangea maderaspatana	Compositae	Η	Mastaru	C&B	LC
163	Grewia flavescens	Malvaceae	Т	Bankajaana	C&B	LC
164	Grewia hirsuta	Malvaceae	S	Nalla Kattelu	C&B	NA
165	Grewia tiliifolia	Malvaceae	Т	Budamara	C&B	NA
166	Grewia villosa	Malvaceae	S	Banta, Chenula	C&B	LC
167	Gymnosporia emarginata	Celastraceae	S	Danthi	C&B	NA
168	Habenaria panigrahiana	Orchidaceae	H	-	С	NA
169	Hackelochloa granularis	Poaceae	Н	Naalipunuku	С	NA

170	Hardwickia binata	Leguminosae	Т	Naarepi C&B		LC
171	Helicteres isora	Malvaceae	S	Gooba thada	C&B	NA
172	Heliotropium indicum	Boraginaceae	Н	Nagadanti	В	NA
173	Heliotropium marifolium	Boraginaceae	Η	-	С	NA
174	Hemidesmus indicus	Apocynaceae	С	Sugandhapala	C&B	NA
175	Heteropogon contortus	Poaceae	Η	Yeddigaddi/ Kes aragaddi	C&B	NA
176	Hibiscus micranthus	Malvaceae	S	Nityamalle	В	NA
177	Holarrhena pubescens	Apocynaceae	S	Kolamukhi	C&B	LC
178	Holoptelea integrifolia	Ulmaceae	Т	Nemalinaara	C&B	NA
179	Huberantha cerasoides	Annonaceae	Т	Chilakadudduga	В	NA
180	Hydrilla verticillata	Hydrocharitacee	AH	Poonaachu	В	LC
181	Hygrophila auriculata	Acanthaceae	AH	Mulla gobbi	В	LC
182	Indigofera linnaei	Leguminosae	Н	Yerrapalleru	C&B	NA
183	Ipomoea aquatica	Convolvulaceae	С	Thootiloora	В	LC
184	Ipomoea carnea	Convolvulaceae	С	Pandiri thooti	В	NA
185	Ipomoea nil	Convolvulaceae	С	Kolivitthulu	В	NA
186	Ipomoea obscura	Convolvulaceae	С	Nallateega	C&B	NA
187	Ipomoea pes-tigridis	Convolvulaceae	С	Mekamadugu	В	NA
188	Ipomoea sagittifolia	Convolvulaceae	С	Purititeega	В	NA
189	Ixora pavetta	Rubiaceae	S	Koravi	C&B	NA
190	Jasminum auriculatum	Oleaceae	L	Adavi teega	C&B	NA
				malli		
191	Jatropha gossypiifolia	Euphorbiaceae	S	Nela amudam	В	LC
192	Justicia glauca	Acanthaceae	Η	-	C&B	NA
193	Justicia vahliana	Acanthaceae	Н	-	C&B	NA
194	Knoxia sumatrensis	Rubiaceae	Η	Kampurodda	C&B	NA
195	Lagerstroemia parviflora	Lythraceae	Т	Chennangi	C&B	NA
196	Lannea coromandelica	Anacardiaceae	Т	Gumpena	C&B	LC
197	Lantana camara	Verbenaceae	S	Seesa kammari	C&B	NA
198	Lepidagathis cristata	Acanthaceae	Η	Nakka pintuka	В	NA
199	Leptopetalum biflorum	Rubiaceae	Η	-	C&B	NA
200	Leucaena leucocephala	Leguminosae	Т	Subabulu	C&B	NA
201	Limonia acidissima	Rutaceae	Т	Velaga	C&B	NA
202	Madhuca longifolia	Sapotaceae	Т	Ірра	C&B	NA
203	Maerua apetala	Capparaceae	Т	Pilli Adugu	В	NA
204	Malvastrum coromandelianum	Malvaceae	Н	-	В	NA
205	Mangifera indica	Anacardiaceae	Т	Mamidi	В	DD
206	Manilkara hexandra	Sapotaceae	Т	Pala chettu	C&B	NA
207	Marsilea quadrifolia	Marsileaceae	AH	Chandamama kura	В	LC
208	Memecylon umbellatum	Melastomataceae	Т	Alli, Peddalli	В	NA
209	Merremia gangetica	Convolvulaceae	H	Elukagemudu	В	LC
210	Mesosphaerum suaveolens	Lamiaceae	H	Danthitulasi	C&B	NA
211	Miliusa tomentosa	Annonaceae	Т	Barredudduga	C&B	NA
212	Mimosa pudica	Leguminosae	H	Attipatti, Mudatha thaamara	В	LC
213	Mitragyna parvifolia	Rubiaceae	Т	Kadamba	C&B	NA
214	Morinda coreia	Rubiaceae	Т	Togaru	C&B	NA
215	Naringi crenulata	Rutaceae	Т	Kukka velaga	В	NA
216	Nelumbo nucifera	Nelumbonaceae	AH	Kamalam	В	NA

217	Nyctanthes arbor-tristis	Oleaceae	Т	Parijatham B		NA
218	Nymphaea nouchali	Nymphaeaceae	AH	Thamara	В	LC
219	Ochna obtusata	Ochnaceae	Т	Raktha sirishamu	C&B	NA
220	Ocimum basilicum	Lamiaceae	Η	Kammagaggiri B aku		NA
221	Ocimum tenuiflorum	Lamiaceae	Н	Thulasi	В	NA
222	Olax scandens	Olacaceae	L	Yelaka Nakkera	C&B	NA
223	Oldenlandia umbellata	Rubiaceae	Н	Chiriveru	В	NA
224	Opuntia stricta	Cactaceae	S	Brahmma jemudu	C&B	LC
225	Orthosiphon rubicundus	Lamiaceae	Н	Podathulasi	C&B	NA
226	Osbeckia zeylanica	Melastomataceae	Н	Burada alli	С	NA
227	Ouret lanata	Amaranthaceae	Н	Kondapindi	В	NA
228	Parthenium hysterophorus	Compositae	Н	Vayyaribhama	C&B	NA
229	Pavonia zeylanica	Malvaceae	Н	Karubenda	В	NA
230	Pergularia daemia	Apocynaceae	С	Dustapa teega	C&B	LC
231	Persicaria glabra	Polygonaceae	AH	Neetiganneru, Burada Gogu	В	LC
232	Phoenix loureiroi	Arecaceae	S	Konda itha	C&B	LC
233	Phoenix sylvestris	Arecaceae	Т	Itha chettu	C&B	NA
234	Phyllanthus amarus	Phyllanthaceae	Н	Nelausiri	В	NA
235	Phyllanthus emblica	Phyllanthaceae	Т	Usirikaya	C&B	LC
236	Phyllanthus maderaspatensis	Phyllanthaceae	Н	Pedda Nela Usiri	C&B	LC
237	Phyllanthus reticulatus	Phyllanthaceae	S	Purugudu	В	LC
238	Phyllanthus virgatus	Phyllanthaceae	Н	Gadhausiri	C&B	NA
239	Pleurolobus gangeticus	Leguminosae	Н	Kolakuponna	C&B	NA
240	Pongamia pinnata	Leguminosae	Т	Kanuga	В	LC
241	Portulaca oleracea	Portulacaceae	Н	Payalaku	В	LC
242	Premna mollissima	Lamiaceae	Т	Kondamanga	C&B	NA
243	Prosopis juliflora	Leguminosae	Т	Sarkar tumma	C&B	NA
244	Pseudarthria viscida	Fabaceae	C	Adavi chikkudu	В	NA
245	Pteris argyraea	Pteridaceae	AH	-	В	NA
246	Pterolobium hexapetalum	Leguminosae	L	Korintha	C&B	NA
247	Pterospermum xylocarpum	Malvaceae	Т	Loluguchettu, Thada	C&B	NA
248	Pulicaria wightiana	Asteraceae	Η	Adavipoddutir ugudu	C&B	NA
249	Pupalia lappacea	Amaranthaceae	Η	Yerra C&B utthareni		LC
250	Rhynchosia rufescens	Leguminosae	C	-	В	NA
251	Rivea hypocrateriformis	Convolvulaceae	L	Bodditeega	C&B	NA
252	Ruellia tuberosa	Acanthaceae	Η	Chitapatakayal a mokka	C&B	NA
253	Saccharum spontaneum	Poaceae	Н	Naagaswaram, Adavicheruku	В	LC
254	Santalum album	Santalaceae	Т	Swethagandham	С	VU
255	Sapindus emarginatus	Sapindaceae	Т	Kunkudu	В	NA
256	Schleichera oleosa	Sapindaceae	Т	Poosuga	C&B	LC
257	Schoenoplectiella articulata	Cyperaceae	Н	-	C&B	LC
258	Schrebera swietenioides	Oleaceae	Т	Magalinga	C&B	NA

259	Scleria lithosperma	Cyperaceae	Н	- C&		NA
260	Scoparia dulcis	Plantaginaceae	Н	Godathulasi	C&B	NA
261	Senegalia chundra	Leguminosae	Т	Sandra	C&B	NA
262	Senegalia torta	Leguminosae	S	Korinta teega	В	NA
263	Senna auriculata	Leguminosae	S	Thangedi	В	NA
264	Senna occidentalis	Leguminosae	Н	Kasinda	В	LC
265	Senna tora	Leguminosae	Η	Thantepu mokka	В	NA
266	Sida acuta	Malvaceae	Η	Katari kaanaaku	В	NA
267	Sida cordifolia	Malvaceae	Н	Badiyalaku	В	NA
268	Solanum virginianum	Solanaceae	Н	Nela vaakudu	В	NA
269	Soymida febrifuga	Meliaceae	Т	Somi	В	NA
270	Spermacoce articularis	Rubiaceae	Н	Madanaku	В	NA
271	Spermacoce pusilla	Rubiaceae	Н	Chukkakaada	В	NA
272	Sphaeranthus indicus	Compositae	Н	Bodasaramu	В	LC
273	Stemodia viscosa	Plantaginaceae	Н	Bodasaramamu	C&B	NA
274	Sterculia urens	Malvaceae	Т	Thapassi	C&B	NA
275	Stereospermum tetragonum	Bignoniaceae	Т	Ummrttha	С	NA
276	Streblus asper	Moraceae	Т	Barraniki	В	LC
277	Striga angustifolia	Orobanchaceae	Н	Jonnamalle	C&B	NA
278	Striga asiatica	Orobanchaceae	Н	Raathibadanika	C&B	NA
279	Strobilanthes pavala	Acanthaceae	Н	-	С	NA
280	Strychnos nux-vomica	Loganiaceae	Т	Mustichettu	C&B	NA
281	Strychnos potatorum	Loganiaceae	Т	Chillanginja chettu	C&B	NA
282	Tacca leontopetaloides	Dioscoreaceae	Н	Ritthakanda	С	LC
283	Tamarindus indica	Leguminosae	Т	Chinta	C&B	LC
284	Tarenna asiatica	Rubiaceae	S	Konda Papidi	C&B	NA
285	Tectona grandis	Lamiaceae	Т	Teku	C&B	NA
286	Tephrosia purpurea	Leguminosae	Н	Vempali	C&B	NA
287	Tephrosia villosa	Leguminosae	Н	Noogu vempalli	В	LC
288	Terminalia anogeissiana	Combretaceae	Т	Sirimanu	C&B	NA
289	Terminalia arjuna	Combretaceae	Т	Tellamaddhi	C&B	NA
290	Terminalia bellirica	Combretaceae	Т	Thani	C&B	LC
291	Terminalia elliptica	Combretaceae	Т	Nallamaddi	В	NA
292	Trianthema portulacastrum	Aizoaceae	Н	Galijeru	C&B	NA
293	Tridax procumbens	Compositae	Н	Gaddichamanthi	В	NA
294	Trigastrotheca pentaphylla	Molluginaceae	Η	Pichichatraku	C	NA
295	Triumfetta rotundifolia	Malvaceae	S	-	C	NA
296	Typha angustifolia	Typhaceae	S	Jammu	В	LC
297	Urochloa ramosa	Poaceae	Н	-	В	LC
298	Vachellia leucophloea	Leguminosae	Т	Tella thumma	C&B	LC
299	Vachellia nilotica	Leguminosae	Т	Nalla thumma	C&B	LC
300	Vanda tessellata	Orchidaceae	ES	Badanika	C&B	LC
301	Ventilago denticulata	Rhamnaceae	L	Erra sanguru	В	NA
302	Viscum articulatum	Santalaceae	S	Jilledu badanika	С	NA
303	Vitex negundo	Lamiaceae	S	Nalla vavilli	C&B	LC
304	Waltheria indica	Malvaceae	S	Nallabenda	C&B	LC
305	Woodfordia fruticosa	Lythraceae	S	Jargi seringi	C&B	LC

306	Wrightia tinctoria	Apocynaceae	Т	Palamaneru	C&B	NA
307	Xanthium strumarium	Compositae	S	Marulamathangi	В	NA
308	Xenostegia tridentata	Convolvulaceae	C	Mududantla	В	NA
309	Xylia xylocarpa	Leguminosae	Т	Bojja	C&B	LC
310	Ziziphus jujuba	Rhamnaceae	Т	Regu	C&B	LC
311	Ziziphus oenopolia	Rhamnaceae	L	Pariki kampa	C&B	LC
312	Ziziphus xylopyrus	Rhamnaceae	Т	Gotti	C&B	NA
313	Zornia gibbosa	Fabaceae	Н	Nelabariki	C&B	NA

*Note: T-Trees; S-Shrubs; H-Herbs; C-Climbers; L-Lianas; AH-Aquatic herb; EP-Epiphyte; C-Core zone; B-Buffer zone; LC-Least Concern; NA-Not Assessed; VU-Vulnerable; NT-Near Threatened



Fig. 12. Vegetation Analysis

Twenty-nine Aquatic species observed in the study area of which, 19 are Angiospems (*Actinoscirpus grossus, Ammannia baccifera, A. multiflora, Aponogeton crispus, Bacopa monnieri, Bergia capensis, Bonnaya ciliate, Canscora heteroclite, Commelina benghalensis, Cyanotis axillaris, Cyperus corymbosus, C.difformis, C. exaltatus, C. mindorensissubsp. pygmaeus, C. mindorensis, C. pangorei, Eleocharis geniculate, Fimbristylis aestivalis, F. argentea, F. quinquangularis, Hydrilla verticillata, Hygrophila auriculata, Nelumbo nucifera, Nymphaea nouchali and Persicaria glabra), three are Pteridophytes (<i>Ceratopteris thalictroides, Marsilea quadrifolia* and *Pteris argyraea*) and *Chara globularis* is Algae.

The most dominant families are Fabaceae represented by 40 species of 31 genera followed by Poaceae 18/14, Malvaceae 18/13, Rubiaceae 16/14, Cyperaceae 15/7, Asteraceae 14/12, Acanthaceae 12/9, Convolvulaceae 11/5, Lamiaceae 9/7, Euphorbiaceae 9/6, Apocyanaceae 8/7, Amaranthaceae 8/6, Phyllanthaceae 8/3, Combretaceae 6/3, Moraceae 6/2, Boraginaceae 5/4, Rutaceae 4/4, Capparaceae & Lythraceae each with 4/3, Rhamnaceae 4/2, Anacardiaceae, Annonaceae, Asparagaceae, Oleaceae, Plantaginaceae and Sapindaceae each with 3/3, Commelinaceae & Molluginaceae each with 3/2, Ebenaceae 3/1, Burseraceae, Celastraceae, Dioscoreaceae, Melastomataceae, Meliaceae, Orchidaceae, Pteridaceae, Santalaceae & Sapotaceae each with 2/2, Arecaceae, Loganiaceae, Nyctaginaceae & Orobanchaceae each with 2 species in 1 generaand 41 families are represented each with one species of one genera.

Top ten families are Fabaceae, Poaceae, Malvaceae, Rubiaceae, Cyperaceae, Asteraceae, Acanthaceae, Convolvulaceae, Lamiaceae and Euphorbiaceae are occupied 52% of the total vegetation structure.



Fig. 13. Top Ten Dominated Families

Fig. 14. Vegetation structure in the proposed project



Phyto-sociological features of trees, shrubs and herbs observed in core & buffer zones:

Core Zone

- A. Trees From Importance Value Index (IVI), it is inferred that for trees the dominant species is *Tectona grandis* is with 34.71 of Importance Value Index followed by *Cleistanthus collinus* (22.97), *Xylia xylocarpa* (20.24) and *Boswellia serrata* (18.73). Simpson's Index value (0.07) and Shannon Wiener Index value (1.24) for trees shows that diversity is more in comparison to species dominance.
- **B.** Shrubs *Butea superb* is identified as the most dominant species with 51.6 of Importance Value Index followed by *Helicteres isora* (42.07), *Waltheria indica* (37.56), *Dioscorea pentaphylla* and *Derris scandens* each with 33.98. Simpson's Index value is 0.13 and the Shannon Wiener Index value is 0.93, which shows that diversity is more.
- **C.** Herbs *Drimia indica* is identified as the most dominant species with 43.85 of Importance Value Index followed by *Andrographis paniculata* and *Cyanthillium albicans* each with 39.21 and *Gloriosa superba* (37.89). Simpson's Index value is 0.13 and the Shannon Wiener Index value is 0.92, which shows that diversity is more.

Core zone					
31.62 x	31.62 m for Trees				
S.No.	Name	IVI	Simpson	Shannon	
1	Tectona grandis	34.71	0.028	-0.13	
2	Albizia odoratissima	8.53	0	-0.04	
3	Givotia moluccana	10.87	0.001	-0.05	
4	Cleistanthus collinus	22.97	0.009	-0.1	
5	Lannea coromandelica	3.74	0	-0.01	
6	Diospyros melanoxylon	17.88	0.004	-0.08	
7	Boswellia serrata	18.73	0.005	-0.08	
8	Dalbergia lanceolaria subsp.paniculata	13.42	0.002	-0.06	
9	Stereospermum tetragonum	8.53	0	-0.04	
10	Lagerstroemia parviflora	15.33	0.002	-0.06	
11	Hardwickia binata	9.7	0.001	-0.04	
12	Acacia leucophloea	11.51	0.001	-0.05	
13	Grewia flavescens	13.22	0.002	-0.06	
14	Hardwickia binata	10.87	0.001	-0.05	
15	Firmiana simplex	12.82	0.001	-0.05	
16	Acacia chundra	17.9	0.004	-0.08	
17	Aegle marmelos	13.22	0.002	-0.06	
18	Chloroxylon swietenia	15.56	0.003	-0.07	
19	Ficus mollis	3.74	0	-0.01	
20	Ochna obtusata	9.19	0	-0.04	
21	Xylia xylocarpa	20.24	0.005	-0.08	
22	Wrightia tinctoria	7.37	0	-0.03	
			0.07	-1.24	

Table 4. Phyto-sociological features observed in core zone

5 x 5 m	l for Shrubs			
1	Butea superba	51.6	0.047	-0.14
2	Cajanus scarabaeoides	25.89	0.006	-0.09
3	Waltheria indica	37.56	0.019	-0.12
4	Phoenix loureiroi	21.5	0.003	-0.07
5	Helicteres isora	42.07	0.025	-0.13
6	Grewia hirsuta	10.99	0	-0.03
7	Dioscorea pentaphylla	33.98	0.014	-0.11
8	Derris scandens	33.98	0.014	-0.11
9	Catunaregam spinosa	17.81	0.002	-0.06
10	Tarenna asiatica	24.63	0.003	-0.07
			0.13	-0.93
1 x 1 m	n for Herbs			
1	Gloriosa superba	37.89	0.02	-0.12
2	Evolvulus nummularius	26.09	0.006	-0.09
3	Cyanthillium albicans	39.21	0.02	-0.12
4	Orthosiphon rubicundus	41.51	0.026	-0.13
5	Andrographis paniculata	39.21	0.02	-0.12
6	Byttneria herbacea	29.55	0.006	-0.09
7	Drimia indica	43.85	0.026	-0.13
8	Scleria lithosperma	24.8	0.004	-0.07
9	Phyllanthus virgatus	17.92	0.002	-0.06

Buffer Zone

- A. Trees From Importance Value Index (IVI), it is inferred that, for trees *Xylia xylocarpa* is the most dominant species with 21.00 of Importance Value Index followed by *Tectona grandis* (16.27), *Lannea coromandelica* (15.85), *Anogeissus latifolia* (14.76), *Schleichera oleosa* and *Cleistanthus collinus* each with (10.3) and *Cleistanthus collinus* (10.36). Simpson's Index value is 0.042 and the Shannon Wiener Index value is 1.47, which shows that diversity is more.
- **B.** Shrubs *Woodfordia fruticosa* is identified as the most dominant species with 35.44 of Importance Value Index followed by *Canthium coromandelicum* (22.86), *Chromolaena odorata* (21.26), *Helicteres isora* (20.55) and *Getonia floribunda* (20.2). Simpson's Index value is 0.07and the Shannon Wiener Index value is 1.19, which shows that diversity is more.
- **C.** Herbs *Hybanthus enneaspermus* is identified as the most dominant species with 25.95 of Importance Value Index followed by *Orthosiphon rubicundus* (25.18), *Oldenlandia umbellate* and *Scleria terrestris* each with 23.84, *Byttneria herbacea* (22.88) and *Lepidagathis cristata* (21.02). Simpson's Index value is 0.07 and the Shannon Wiener Index value is 1.18, which shows that diversity is more.

This infers the buffer zone of woody species has more diversity when compared to other life forms of plants. Most of the natural growth recorded is of coppice in nature.
Buffer zone								
31.62	x 31.62 m for Trees							
1	Xylia xylocarpa	21	0.01	-0.1				
2	Lannea coromandelica	15.85	0.005	-0.08				
3	Anogeissus latifolia	14.76	0.004	-0.07				
4	Madhuca longifolia var. latifolia	9.62	0.001	-0.05				
5	Mitragyna parvifolia	7	0	-0.03				
6	Schleichera oleosa	10.37	0.001	-0.05				
7	Terminalia alata	9.17	0.001	-0.05				
8	Miliusa tomentosa	10.08	0.001	-0.05				
9	Hardwickia binata	8.71	0.001	-0.04				
10	Cleistanthus collinus	10.36	0.001	-0.05				
11	Boswellia serrata	8.01	0.001	-0.04				
12	Buchanania cochinchinensis	8.26	0.001	-0.04				
13	Soymida febrifuga	7	0	-0.03				
14	Bombax ceiba	9.17	0.001	-0.05				
15	Albizia odoratissima	8.51	0.001	-0.04				
16	Givotia moluccana	8.09	0.001	-0.04				
17	Diospyros melanoxylon	7	0	-0.03				
18	Bridelia retusa	6.33	0	-0.03				
19	Dalbergia lanceolaria subsp.paniculata	9.02	0.001	-0.05				
20	Careya arborea	7.5	0.001	-0.04				
21	Lagerstroemia parviflora	9.02	0.001	-0.05				
22	Tectona grandis	16.27	0.005	-0.08				
23	Pterospermum xylocarpum	6.33	0	-0.03				
24	Terminalia bellirica	4.49	0	-0.02				
25	Grewia tiliifolia	6.92	0	-0.03				
26	Butea monosperma	4.49	0	-0.02				
27	Gardenia latifolia	6.33	0	-0.03				
28	Chloroxylon swietenia	4.49	0	-0.02				
29	Albizia amara	5.99	0	-0.03				
30	Bauhinia racemosa	5.99	0	-0.03				
31	Aegle marmelos Wrightig tingtonig	4.48	0	-0.02				
32	wrightia thictoria	9.75	0.001	-0.04				
33	Acadia laucophloca	/.49	0	-0.03				
34	Acucia leucophioea	4.40	0	-0.02				
35	Luphorbia nivulta Naringi granulata	4.40	0	-0.02				
30		3.23	0 042	-0.01				
5 7 5 7	n for Shruhs		0.042	-1.4/				
1	Woodfordia fruticosa		0.025	-0 19				
- I 	Tarenna asiatica	18 00	0.025	-0.13				
2	Ixora pavetta	15.09	0.004	-0.07				
	Helicteres isora	20.55	0.005	-0.02				
	Canthium coromandelicum	20.00	0.005	-0.08				
6	Getonia floribunda	20.2	0.006	-0.00				

Table 5. Phyto-sociological features observed in core zone

7	Chromolaena odorata	21.26	0.007	-0.09
8	Catunaregam spinosa	19.14	0.005	-0.08
9	Erythroxylum monogynum	18.09	0.004	-0.08
10	Grewia hirsuta	12.3	0.001	-0.05
11	Ziziphus oenopolia	15.72	0.003	-0.07
12	Carissa carandas	12.3	0.001	-0.05
13	Butea superba	12.3	0.001	-0.05
14	Olax scandens	11.31	0.001	-0.04
15	Acalypha alnifolia	6.46	0	-0.02
16	Dichrostachys cinerea	8.89	0	-0.03
17	Flacourtia indica	6.46	0	-0.02
18	Phoenix loureiroi	6.46	0	-0.02
19	Triumfetta rotundifolia	16.16	0.001	-0.05
			0.07	-1.19
1 x 1 r	n for Herbs			
1	Byttneria herbacea	22.88	0.008	-0.09
2	Orthosiphon rubicundus	25.18	0.01	-0.1
3	Andrographis paniculata	15.6	0.002	-0.06
4	Spermacoce articularis	18.51	0.004	-0.08
5	Sida cordifolia	17.06	0.003	-0.07
6	Scleria lithosperma	23.84	0.008	-0.09
7	Oldenlandia umbellata	23.84	0.008	-0.09
8	Phyllanthus virgatus	17.74	0.004	-0.08
9	Hybanthus enneaspermus	25.95	0.011	-0.1
10	Lepidagathis cristata	21.02	0.006	-0.09
11	Desmodium gangeticum	16.26	0.003	-0.07
12	Fimbristylis ovata	20.91	0.003	-0.07
13	Evolvulus nummularius	11.5	0.001	-0.04
14	Eragrostis viscosa	8.36	0	-0.03
15	Elytraria acaulis	11.5	0.001	-0.04
16	Cyanthillium albicans	8.36	0	-0.03
17	Alysicarpus monilifer	11.5	0.001	-0.04
			0.07	-1.18

IUCN Red list species

During primary data collection, identified *Chloroxylon swietenia*, *Cleistanthus collinus*, *Curcuma pseudomontana*, *Cycas rumphii* (Planted in core zone) and *Santalum album* which are falls under Vulnerable (Vu) category and *Albizia thompsonii* listed in Near Threatened category as per IUCN Red list.

Seed collection and rising of saplings may also be taken up by SCCL in their nurseries and planted in open areas to restore the ecosystem. *Cycas rumphii* was planted 30 years ago in Project Office of GK-OCP campus.

Density enrichment plantations by Forest Department within the 10 Kms radius:

The forest department has raised density enrichment plantations in a total of 234 Hectares, comprising 5 forest beats within the radius of 10 Kms, which is being home range for varied wildlife existing and moving in the adjoining forests outside the mine boundaries.

Also, the department has developed a 15 acres of grass plot in Tippanapalli forest beat as a fodder/ feeding ground for the ungulates present in the forest areas within the 10 Kms radius.

S.No.	Beat		Plantation	in Ha	Grass land		and
		На	Latitude	Longitude	На	Latitude	Longitude
1	Tungaram	30	17.44781	80.61903	0		
		20	17.4386	80.61918			
		20	17.43909	80.62068			
		10	17.4454	80.62151			
		5	17.44800	80.61940			
		14	17.43915	80.62355			
	Sub-Total:	99			0		
2	Tippanapalli	20	17.4364	80.62889	15	17.43769	80.64544
		25	17.4296	80.62501			
		25	17.4287	80.61866			
		20	17.42656	80.63245			
		25	17.42041	80.62945			
	Sub-Total:	115			15		
3	Penagadapa	20	17.42395	80.68048	0		
	Sub-Total:	20			0		
4	Ramavaram	0			0		
	Sub-Total:	0			0		
5	Gareebpeta	0			0		
	Sub-Total:	0			0		
	Grand Total:	234			15		

Table 6. Total area of Beat wise plantations by forest department

The proposed core zone of the project is falls in Ramavaram reserve forest and Buffer zone is occupied by Five RFs namely falling in Ramavaram RF, Chatakonda RF, Penagadapa RF, Mulakalapalli RF and Paloncha East RF of Kothagudem and Paloncha divisions.

There are no National Parks, Biosphere Reserve, Tiger Reserves and Elephant Reserve/Corridor in the study area.

Density enrichment plantations by SCCL within the 10 Kms radius:

A total of 1106 Hectares of (block, OB, and avenue) plantations have been raised by SCCL since 1984 in this area during ecological restoration processes. These restoration practices are highly upsurging in reversing the habitat degradation happened in and around the mine areas.

During the survey, it is observed that all the plantations are helping to restore organisms and their interactions with one another and with the physical environment. Observations of varied faunal groups shows the success of the ecological restoration practices of the mines, which focusses on the processes such as perseverance of species through natural recruitment and survival, functioning food webs, system-wide nutrient conservation via relationships among plants, and animals.

Year	Location	Name of	Area	Lat & Long
ot		the	in Ha	
raising		planted		
1984	Cashew plantation	Cashew	27	N 17.5267 E 80.6424
1999	Block pl' at CRP Camp	EC	9	N 17.5263 E 80.6311
1999	Garemellapadu	EC	11	N 17.5267 E 80.6424
2000	7 Shaft dispensary	EC	14	N 17.4725 E 80.6656
2000	Garemellapadu	EC	5	N 17.5191 E 80.6442
2001	Garemellapadu	EC	5	
2003	3 Incline	EC	12	N 17.5074 E 80.6479
2003	Hemachandrapuram	EC	10	N 17.5535 E 80.5916
2003	Garemellapadu	EC	19	N 17.5209 E 80.6467
2004	Rly.siding	EC	7	N 17.5050 E 80.6518
2004	St. Josephs	EC&M	7	N 17.4983 E 80.6510
2004	VK Timberyard	EC	5	N 17.4692 E 80.6668
2004	Coal Testing Lab	EC	8	N 17.5061 E 80.6528
2005	GPF Nursery	EC	25	N 17.5345 E 80.6314
2005	GKOCP	М	10	N 17.5345 E 80.6315
2005	5 & 7 Shaft	EC,M&B	35	N 17.4830 E 80.6647
2005	VK compound wall	EC&M	7	N 17.4783 E 80.6729
2006	Vanamanagar	EC&M	5	N 17.4702 E 80.6682
2006	Goutampur	М	4	N 17.4815 E 80.6446
2006	5Shaft	М	10	N 17.4968 E 80.6692
2006	Gadalamadugu	EC	13	N 17.5423 E 80.5562
2006	GKOCP	М	10	N 17.5423 E 80.5563
2007	Tilaknagar	EC	5	N 17.4802 E 80.6474
2007	Goutampur	EC&M	3	N 17.4828 E 80.6463
2007	Behind 3 Incline	EC	5	N 17.5100 E 80.6414
2007	7 Incline	EC&M	5	N 17.4748 E 80.6678
2007	7 Incline (S)	EC&M	8	N 17.4838 E 80.6719
2007	2 Incline	EC	4	N 17.5300 E 80.6382
2007	Tellavagu	М	6	N 17.4931 E 80.6850
2008	Subsidence panel 1 & 1A	М	6	N 17.4922 E 80.6625
2009	Behind VK-7 Incline Mine-RF	EC	9	N17.5770 E 80.6379
2009	Evacuated Colony of VK-7 Incline	EC	7	N 17.4761 E 80.6680

Table 7. Kothagudem Block plantations from 1984-2020 by SCCL

2009	Gaps in cashew pl' near Dhanbad and ITI	EC	24	N 17.5223 E 80.6411
2009	Gaps in cashew pl' near Christian graveyards 2 incline	EC	6	N 17.4923 E 80.6584
2009	Gaps in cashew pl' near B CF	EC	8	N 17.5147 E 80.6439
2009	Behind Central Nursery, KGM	Μ	9	N 17.5770 E 80.6378
2010	GKOCP-Block	EC&M	8	N 17.4466 E 80.6609
2011	RF area in KGM near Deport.	М	15	N 17.5724 E80.6327
2012	GKOCP Vacant patches	М	4	N 17.4622 E80.65163
2012	VK-7 Incline Vacant patches	Μ	4	N 17.4816 E80.67233
2013	Gaps in Cashew plantations at Garimellapadu	Μ	33	N 17.3130 E80.3835
2014	VK-7 Incline	EU	10	N 17.2848 E80.4014
2016	RF plantation near Gaddalamadugu	Μ	22	N 17.3301 E80.3830
2016	RF plantation near 5B shaft	М	42	N 17.3012 E80.4018
2016	Euc.clonal planatation near CRP camp, Ramavaram	EC	2	N 17.5283 E80.6309
2016	Vacant patches in writer basthi. KGM/Corp	М	2	N 17.5283 E80.6310
2017	RF plantation at Gaddalamadugu	EC & M	50	N 17.4947 E.80.657457
	Gap plantation from writer basthi to main hospital area, Kothagudem	Μ	10	N 17.5500 E80.608049
2019	Garimellapadu	Μ	2	N 17.5272 E.80.648357
	Total		567.00	

(EC- Eucalyptus; B – Bamboo; M – Mixed/Miscellaneous)

Table 8. Kothagudem Avenue plantations by SCCL

Year of raising	Location	Name of the Species planted	Area in Ha	Lat & Long
2001	Goutampur	Μ	2	N 17.4809 E 80.6478
2002	Rudrampur	Μ	5	N 17.4984 E 80.6527
2003	GKOCP, Writer basthi, Rudrampur colony	М	13	N 17.4555 E 80.6283
2004	Rudrampur,5shaft	М	5	N 17.4984 E 80.6527
2005	5&7 shaft	М	5	N 17.4842 E 80.6679
2008	Kothagudem area	Μ	2	N 17.4988 E 80.6530
2009	Gaps in Writer basthi, KGM	Μ	3	N 17.5467 E 80.6156
2010	GKOCP	Μ	2	N 17.4466 E 80.6591
2014	PVK-Guest House	М	1	N 17.2911 E80.3929
	Total		38.00	

Year of raising	Location	Name of the Species planted	Area in Ha	Lat & Long
1999	GKOCP, OB	Μ	2	N 17.4457 E 80.6456
2001	GKOCP, OB	Μ	10	N 17.4495 E 80.6441
2002	GKOCP, OB	Μ	23	N 17.4578 E 80.6285
2003	GKOCP, OB	Μ	52	N 17.4558 E 80.6329
2004	GKOCP, OB	Μ	65	N 17.4585 E 80.6294
2005	GKOCP, OB	Μ	48	N 17.4640 E 80.6298
2006	GKOCP, OB	Μ	31	N 17.4618 E 80.6328
2007	GKOCP, OB	Μ	7	N 17.4600 E 80.6368
2008	GKOCP, OB	Μ	8	N 17.2086 E 80.8122
2009	GKOCP-West Dump-RF, OB	Μ	44	N 17.4618 E 80.6328
2010	GKOCP-OB	Μ	28	N 17.4466 E 80.6591
2011	GKOCP, OB	Μ	19	N17.44881 E80.65703
2012	GKOCP OB dumps	Μ	25	N17.44663 E80.65681
2013	GKOCP, OB	Μ	33	N17.2643 E80.3919
2014	GKOCP, OB	Μ	12	N17.2871 E80.3949
2015	GKOCP, OB	Μ	12	N17.2718 E80.3815
2016	GKOCP, OB	Μ	24	N17.2871 E80.3949
2017	GKOCP, OB	Μ	12	N17.4519 E80.6466
2018	GKOCP, OB	EC & M	46	N17.4511 E80.6478
	Total for Kothagudem		501.00	

Table 9. GKOCP OB plantations by SCCL

Planting suitable vegetation can help to re-establish habitats relatively rapidly after mining disturbance. This may benefit varied biodiversity/wildlife of the area.

List of mixed/miscellaneous species planted by SCCL:

- 1. Hardwikia binata
- 2. Dendrocalamus strictus
- 3. Ficus religiosa
- 4. Pterocarpus santalinus
- 5. Azadirachta indica
- 6. Limonia acidissima
- 7. Ficus bengalensis
- 8. Aegle marmelos
- 9. Mitragyna parvifolia
- 10. Dalbergia latifolia
- 11. Pterocarpus marsupium
- 12. Emblica officinalis
- 13. Syzygium cumini
- 14. Alstonia scholaris
- 15. Albizzia procera

- 16. Annona squamosa
- 17. Terminalia
 - bellarica
- 18. Spathodea campanulata
- 19. Pongamia pinnata
- 20. Ficus mollis
- 21. Lagerstroemia parviflora
- 22. Bauhinia purpurea
- 23. Millingtonia
- harrtensis
- 24. Madhuka indica
- 25. Ficus carica
- 26. Tamarindus indica
- 27. Cassia fistula
- 28. Anthocephalus kadamba
- 29. Pithecelobium dulci

- 30. Sterculia urens
- 31. Tectonia grandis
- 32. Dalbergia sissoo
- 33. Cassia siamea
- 34. Adina cordifolia
- 35. Putranjeeva rasuburgi
- 36. Bombax ceiba
- 37. Mimusops elangii
- 38. Terminalia chebula
- 39. Peltophorum ferruginum
- 40. Albizzia lebbek
- 41. Couravpita guinensis
- 42. Holoptelea integrifolia

Fauna Diversity:

Mammals: In the study area, a total of **19 species** of mammals distributed in 16 Genera belonging to 6 orders and 13 families were recorded. The species include Wild Boar, Common Mongoose, Greater Indian fruit bat, Rhesus Macaque, Bonnet Macaque, Indian Palm Squirrel, Little Indian Field Mouse, Indian hare are common in their occurrence.

S.No.	Common Name	Scientific Name	Family	Order	IUCN Status	IWPA Schedule
1	Nilgai	Boselaphus tragocamelus	Bovidae	Artiodactyla	LC	III
2	Indian Spotted Deer	Axis axis	Cervidae	Artiodactyla	LC	III
3	Indian Sambar	Rusa unicolor	Cervidae	Artiodactyla	VU	III
4	Jungle Cat	Felis chaus	Felidae	Artiodactyla	LC	II
5	Common Grey Mongoose	Urva edwardsii	Herpestidae	Artiodactyla	LC	IV
6	Wild Boar	Sus scrofa	Suidae	Artiodactyla	LC	III
7	Indian Spotted Chevrotain	Moschiola indica	Tragulidae	Artiodactyla	LC	NL
8	Indian Fox	Vulpes bengalensis	Canidae	Carnivora	LC	II
9	Indian Flying Fox	Pteropus medius	Pteropodidae	Chiroptera	LC	IV
10	Indian Hare	Lupus nigricollis	Leporidae	Lagomorpha	LC	IV
11	Rhesus Macaque	Macaca mulatta	Cercopithecidae	Primates	LC	II
12	Bonnet Macaque	Macaca radiata	Cercopithecidae	Primates	VU	II
13	Hanuman Langur	Semnopithecus entellus	Cercopithecidae	Primates	LC	II
14	Indian Crested Porcupine	Hystrix indica	Hystricidae	Rodentia	LC	IV
15	Lesser Bandicoot Rat	Bandicota bengalensis	Muridae	Rodentia	LC	IV
16	Greater Bandicoot Rat	Bandicota indica	Muridae	Rodentia	LC	IV
17	Little Indian Field Mouse	Mus booduga	Muridae	Rodentia	LC	IV
18	House Mouse	Mus musculus	Muridae	Rodentia	LC	IV
19	Indian Palm Squirrel	Funambulus palmarum	Sciuridae	Rodentia	LC	IV

Table 10. List of Mammals recorded from the proposed study area

Herpetofauna: It is represented with **26 species** belonging to 23 genera, 3 orders and 13 families; it includes reptiles and amphibians consisting of two major classes namely Squamata and Anura, of which 20 species of reptiles belonging to orders Squamata and Testudines and 6 species of amphibians belonging to order Anura.

S.No.	Common Name	Scientific Name	Family	Order	Class	IUCN Status	IWPA Schedule
1	Asian Grass Frog	Fejervarya limnocharis	Dicroglossidae	Anura	Amphibia	LC	IV
2	Common Tree Frog	Polypedates leucomystax	Rhacophoridae	Anura	Amphibia	LC	IV
3	Asian Common Toad	Duttaphrynus melanostictus	Bufonidae	Anura	Amphibia	LC	IV
4	Indian Skipper Frog	Euphlyctis cyanophlyctis	Dicroglossidae	Anura	Amphibia	LC	IV
5	Green Pond Frog	Euphlyctis hexadactylus	Dicroglossidae	Anura	Amphibia	LC	IV
6	Indian Bullfrog	Hoplobatrach us tigerinus	Dicroglossidae	Anura	Amphibia	LC	IV
7	Common Garden Lizard	Calotes versicolor	Agamidae	Squamata	Reptilia	LC	IV
8	Forest Blood Sucker	Monilesaurus rouxii	Agamidae	Squamata	Reptilia	LC	NL
9	Pondichéry Fan- throated Lizard	Sitana ponticeriana	Agamidae	Squamata	Reptilia	LC	NL
10	Red sand boa	Eryx johnii	Boidae	Squamata	Reptilia	NT	IV
11	Rough- scaled Sand Boa	Gongylophis conicus	Boidae	Squamata	Reptilia	NT	IV
12	Indian Chameleon	Chamaeleo zeylanicus	Chamaeleonidae	Squamata	Reptilia	LC	II
13	Bronze- backed Tree Snake	Dendrelaphis tristis	Colubridae	Squamata	Reptilia	LC	IV
14	Checkered Keelback	Fowlea piscator	Colubridae	Squamata	Reptilia	LC	II
15	Indian Wolf Snake	Lycodon aulicus	Colubridae	Squamata	Reptilia	LC	IV
16	Common Kukri Snake	Oligodon arnensis	Colubridae	Squamata	Reptilia	LC	IV

 Table 11. List of Herpetofauna recorded from the proposed study area

17	Green Vine Snake	Oxybelis fulgidus	Colubridae	Squamata	Reptilia	LC	IV
18	Indian Rat Snake	Ptyas mucosa	Colubridae	Squamata	Reptilia	LC	IV
19	Indian Cobra	Naja naja	Elapidae	Squamata	Reptilia	LC	II
20	Brooke's House Gecko	Hemidactylus brookii	Gekkonidae	Squamata	Reptilia	LC	NL
21	Common House Gecko	Hemidactylus frenatus	Gekkonidae	Squamata	Reptilia	LC	NL
22	Termite Hill Gecko	Hemiductylus triedrus	Gekkonidae	Squamata	Reptilia	LC	NL
23	Keeled Indian Mabuya	Eutropis carinata	Scincidae	Squamata	Reptilia	LC	NL
24	Common Indian Monitor	Varanus bengalensis	Varanidae	Squamata	Reptilia	NT	Ι
25	Russell's Viper	Daboia russelii	Viperidae	Squamata	Reptilia	LC	II
26	Indian Flapshell Turtle	Lissemys punctata	Trionychidae	Testudines	Reptilia	VU	IV

Avifauna: A total of **86 bird species** belongs to 70 genera, 18 orders and 42 families were recorded in core and buffer zone areas. Among them, order Passeriformes represented with high number of species (37 species distributed in 28 genera of 19 families), followed by Pelecaniformes (8/7/3), Columbiformes & Gruiformes each with 5/4/1, Accipitriformes (4/4/1), Coraciiformes (4/3/3), Galliformes (4/3/1), Apodiformes (3/2/1), Bucerotiformes & Charadriiformes each with 2/2/2, Anseriformes, Ciconiiformes & Cuculiformes each with 2/2/1, Psittaciformes (2/1/1), Falconiformes, Piciformes, Podicipediformes and Strigiformes are represented with 1 species belong to 1 genera of 1 family respectively

Table 12. List of Avifauna	recorded from the	proposed study area
----------------------------	-------------------	---------------------

S.No.	Common Name	Scientific Name	Family	Order	IUCN Status	IWPA Schedule
1	Shikra	Accipiter badius	Accipitridae	Accipitriformes	LC	IV
2	White-eyed Buzzard	Butastur teesa	Accipitridae	Accipitriformes	LC	IV
3	Black-winged Kite	Elanus caeruleus	Accipitridae	Accipitriformes	LC	IV
4	Black Kite	Milvus migrans	Accipitridae	Accipitriformes	LC	IV
5	Lesser Whistling Duck	Dendrocygna javanica	Anatidae	Anseriformes	LC	IV
6	Cotton Teal	Nettapus coromandelianus	Anatidae	Anseriformes	LC	IV

7	Common Swift	Apus apus	Apodidae	Apodiformes	LC	IV
8	House Swift	Apus nipalesis	Apodidae	Apodiformes	LC	IV
9	Asian Palm Swift	Cypsiurus balasiensis	Apodidae	Apodiformes	LC	IV
10	Indian Grey Hornbill	Ocyceros birostris	Bucerotidae Bucerotiformes		LC	IV
11	Eurasian Hoopoe	Upupa epops	Upupidae	Bucerotiformes	LC	IV
12	Red-wattled Lapwing	Vanellus indicus	Charadriidae	Charadriiformes	LC	IV
13	Bronze- winged Jacana	Metopidius indicus	Jacanidae	Charadriiformes	LC	IV
14	Asian Openbill	Anastomus oscitans	Ciconiidae	Ciconiiformes	LC	IV
15	Painted Stork	Mycteria leucocephala	Ciconiidae	Ciconiiformes	NT	IV
16	Rock Pigeon	Columba livia	Columbidae	Columbiformes	LC	IV
17	Spotted Dove	Spilopelia chinensis	Columbidae	Columbiformes	LC	IV
18	Laughing Dove	Spilopelia senegalensis	Columbidae	Columbiformes	LC	IV
19	Oriental Turtle Dove	Streptopelia orientalis	Columbidae	Columbiformes	LC	IV
20	Yellow-footed Green Pigeon	Treron phoenicoptera	Columbidae	Columbiformes	LC	IV
21	White- throated Kingfisher	Halcyon smyrnensis	Alcedinidae	Coraciiformes	LC	IV
22	Little Green Bee- eater	Merops orientalis	Meropidae	Coraciiformes	LC	IV
23	Blue-tailed Bee- eater	Merops philippinus	Meropidae	Coraciiformes	LC	IV
24	Indian Roller	Coracias benghalensis	Coraciidae	Coraciiformes	LC	IV
25	Pied Crested Cuckoo	Clamator jacobinus	Cuculidae	Cuculiformes	LC	IV
26	Asian Koel	Eudynamys scolopaceus	Cuculidae	Cuculiformes	LC	IV
27	Common Kestrel	Falco tinnunculus	Falconidae	Falconiformes	LC	IV
28	Rain Quail	Coturnix coromandelica	Phasianidae	Galliformes	LC	IV
29	Common Quail	Coturnix coturnix	Phasianidae	Galliformes	LC	IV
30	Grey Francolin	Ortygornis pondicerianus	Phasianidae	Galliformes	LC	IV
31	Indian Peafowl	Pavo cristatus	Phasianidae	Galliformes	LC	Ι
32	White- breasted Waterhen	Amaurornis phoenicurus	Rallidae	Gruiformes	LC	IV

33	Common Coot	Fulica atra	Rallidae	Gruiformes	LC	IV
34	Watercock	Gallicrex cinerea	Rallidae	Gruiformes	LC	IV
35	Common Moorhen	Gallinula chloropus	Rallidae	Gruiformes	LC	IV
36	Purple Swamphen	Porphyrio porphyrio	Rallidae	Gruiformes	LC	IV
37	Paddyfield Warbler	Acrocephalus agricola	Acrocephalidae	Passeriformes	LC	IV
38	Booted Warbler	Iduna caligata	Acrocephalidae	Passeriformes	LC	IV
39	Common Iora	Aegithina tiphia	Aegithinidae	Passeriformes	LC	IV
40	Ashy- crowned Sparrow Lark	Eremopterix griseus	Alaudidae	Passeriformes	LC	IV
41	Ashy Woodswallow	Artamus fuscus	Artamidae	Passeriformes	LC	IV
42	Small Minivet	Pericrocotus cinnamomeus	Campephagidae	Passeriformes	LC	IV
43	Common Tailorbird	Orthotomus sutorius	Cisticolidae	Passeriformes	LC	IV
44	Plain Prinia	Prinia inornata	Cisticolidae	Passeriformes	LC	IV
45	Ashy Prinia	Prinia socialis	Cisticolidae	Passeriformes	LC	IV
46	Jungle Prinia	Prinia sylvatica	Cisticolidae	Passeriformes	LC	IV
47	Large-billed Crow	Corvus macrorhynchos	Corvidae	Passeriformes	LC	IV
48	House Crow	Corvus splendens	Corvidae	Passeriformes	LC	V
49	Rufous Treepie	Dendrocitta vagabunda	Corvidae	Passeriformes	LC	IV
50	Tickell's Flowerpecker	Dicaeum erythrorhynchos	Dicaeidae	Passeriformes	LC	IV
51	White-bellied Drongo	Dicrurus caerulescens	Dicruridae	Passeriformes	LC	IV
52	Black Drongo	Dicrurus macrocercus	Dicruridae	Passeriformes	LC	IV
53	Indian Silverbill	Euodice malabarica	Estrildidae	Passeriformes	LC	IV
54	Scaly-breasted Munia	Lonchura punctulata	Estrildidae	Passeriformes	LC	IV
55	Red-rumped Swallow	Cecropis daurica	Hirundinidae	Passeriformes	LC	IV
56	Barn Swallow	Hirundo rustica	Hirundinidae	Passeriformes	LC	IV
57	Wire-tailed Swallow	Hirundo smithii	Hirundinidae	Passeriformes	LC	IV
58	Long-tailed Shrike	Lanius schach	Laniidae	Passeriformes	LC	IV
59	Common Babbler	Argya caudata	Leiothrichidae	Passeriformes	LC	IV
60	Large Grey Babbler	Argya malcolmi	Leiothrichidae	Passeriformes	LC	IV
61	Indian Paradise flycatcher	Terpsiphone paradisi	Monarchidae	Passeriformes	LC	IV

62	Paddyfield Pipit	Anthus rufulus	Motacillidae	Passeriformes	LC	IV
63	White Wagtail	Motacilla alba	Motacillidae	Passeriformes	LC	IV
64	White-browed Wagtail	Motacilla maderaspatensis	Motacillidae	Passeriformes	LC	IV
65	Purple Sunbird	Cinnyris asiaticus	Nectariniidae	Passeriformes	LC	IV
66	Purple- rumped Sunbird	Leptocoma zeylonica	Nectariniidae	Passeriformes	LC	IV
67	Yellow- throated Sparrow	Gymnoris xanthocollis	Passeridae	Passeriformes	LC	IV
68	House Sparrow	Passer domesticus	Passeridae	Passeriformes	LC	IV
69	Red-vented Bulbul	Pycnonotus cafer	Pycnonotidae	Passeriformes	LC	IV
70	Red- whiskered Bulbul	Pycnonotus jocosus	Pycnonotidae	Passeriformes	LC	IV
71	White-browed Bulbul	Pycnonotus luteolus	Pycnonotidae	Passeriformes	LC	IV
72	Common Myna	Acridotheres tristis	Sturnidae	Passeriformes	LC	IV
73	Brahminy Myna	Sturnia pagodarum	Sturnidae	Passeriformes	LC	IV
74	Grey Heron	Ardea cinerea	Ardeidae	Pelecaniformes	LC	IV
75	Purple Heron	Ardea purpurea	Ardeidae	Pelecaniformes	LC	IV
76	Indian Pond Heron	Ardeola grayii	Ardeidae	Pelecaniformes	LC	IV
77	Cattle Egret	Bubulcus ibis	Ardeidae	Pelecaniformes	LC	IV
78	Little Egret	Egretta garzetta	Ardeidae	Pelecaniformes	LC	IV
79	Little Cormorant	Microcarbo niger	Phalacrocoracidae	Pelecaniformes	LC	IV
80	Indian Cormorant	Phalacrocorax fuscicollis	Phalacrocoracidae	Pelecaniformes	LC	IV
81	Red-naped Ibis	Pseudibis papillosa	Threskiornithidae	Pelecaniformes	LC	IV
82	Coppersmith Barbet	Psilopogon haemacephalus	Megalaimidae	Piciformes	LC	IV
83	Little Grebe	Tachybaptus ruficollis	Podicipedidae	Podicipediformes	LC	IV
84	Plum-headed Parakeet	Psittacula cyanocephala	Psittacidae	Psittaciformes	LC	IV
85	Rose-ringed Parakeet	Psittacula krameri	Psittacidae	Psittaciformes	LC	IV
86	Spotted Owlet	Athene brama	Strigidae	Strigiformes	LC	IV

PART	MIN	ΜΑΧ	MEAN (n=2)	SD	Cum Richness
Α	6	8	7.00	0.63	25
В	4	10	7.17	2.56	25
С	3	7	4.50	1.38	14

The total numbers of individuals recorded in all the points are analysed and tabulated.

From the table 12, it is found that the highest number of (20max & 5 min) individuals was recorded in the Part - B (Table 12). The overall density of birds in this point is 49.2 hac.

PART	MIN	ΜΑΧ	MEAN (n=2)	SD	DENSITY in hectare
Α	6	12	8	2.2	40.7
В	5	20	9.7	5.5	49.2
C	3	8	5.7	1.9	28.8

The reason behind the highest record of the individuals in the Part - B is due to the habitat type. The Part has shrub jungle with thick canopies, water sources and less anthropogenic and cattle movements. Hence it shows a high number of species and as well as the highest record of individuals in this part. The minimum number of birds recorded was in Part - C. This is due to rocky terrain and located roadside.

In most of the cases the birds prefer a good habitat for the survival, fulfil their food requirements. So here, on seeing the density of the species we can conclude that the reason for the abundant density in this part is due to their characteristic feature of the habitats.

Bees, Dragonflies, Spiders and Butterflies (Invertebrates): A total of **46 species of Invertebrates**, of which Butterflies under the order Lepidoptera represented with 31 species belong to 26 genera of 5 families. Dragonflies under the order Odonata (9 species belong to 7 genera of 4 families). Hymenoptera is with 4 species belongs to 2 genera of one family. Spiders under the order Araneae represented with 1 species belong to 1 genus of Lycosidae family.

Table 15. List of Invertebrates recorded from the proposed study area

S. No.	Common Name	Scientific Name	Family	Order	Class	IUCN Status	IWPA Schedule
1	Common Funnel Web Spider	Hippasa agelenoides	Lycosidae	Araneae	Arachnida	NA	NL
2	Indian Honeybee	<i>Apis cerana</i> subsp. <i>indica</i>	Apidae	Hymenoptera	Insecta	NA	NL
3	Honeybee	Apis mellifera	Apidae	Hymenoptera	Insecta	NA	NL

4	Violet Carpenter Bee	Xylocopa violacea	Apidae	Hymenoptera	Insecta	NA	NL
5	Carpenter Bee	Xylocopa tenuiscapa	Apidae	Hymenoptera	Insecta	NA	NL
6	Paper Wasp	Ropalidia marginata	Vespidae	Hymenoptera	Insecta	NA	NL
7	Common Bush Hopper	Ampittia dioscorides	Hesperiidae	Lepidoptera	Insecta	NA	NL
8	Common Awl	Hasora badra	Hesperiidae	Lepidoptera	Insecta	NA	NL
9	Indian Grizzled Skipper	Spialia galba	Hesperiidae	Lepidoptera	Insecta	NA	NL
10	Dark Palm Dart	Telicota ancilla	Hesperiidae	Lepidoptera	Insecta	NA	NL
11	Angled Pierrot	Caleta decidia	Lycaenidae	Lepidoptera	Insecta	NA	NL
12	Common Pierrot	Castalius rosimon	Lycaenidae	Lepidoptera	Insecta	NA	NL
13	Lime Blue	Chilades lajus	Lycaenidae	Lepidoptera	Insecta	NA	NL
14	Grass Jewel	Freyeria trochylus	Lycaenidae	Lepidoptera	Insecta	NA	NL
15	Pale Grass Blue	Pseudozizeeria maha	Lycaenidae	Lepidoptera	Insecta	NA	NL
16	Dark Grass Blue	Zizeeria karsandra	Lycaenidae	Lepidoptera	Insecta	NA	NL
17	Blue Grass	Zizina labradus	Lycaenidae	Lepidoptera	Insecta	NA	NL
18	Tiny Grass Blue	Zizula hylax	Lycaenidae	Lepidoptera	Insecta	NA	NL
19	Tawny Coster	Acraea terpsicore	Nymphalidae	Lepidoptera	Insecta	NA	NL
20	Common Castor	Ariadne merione	Nymphalidae	Lepidoptera	Insecta	NA	NL
21	Plain Tiger	Danaus chrysippus	Nymphalidae	Lepidoptera	Insecta	NA	NL
22	Common Tiger	Danaus genutia	Nymphalidae	Lepidoptera	Insecta	NA	NL
23	Common crow	Euploea core	Nymphalidae	Lepidoptera	Insecta	LC	NL
24	Danaid Eggfly	Hypolimnas misippus	Nymphalidae	Lepidoptera	Insecta	NA	NL
25	Peacock Pansy	Junonia almana	Nymphalidae	Lepidoptera	Insecta	NA	NL
26	Yellow Pansy	Junonia hierta	Nymphalidae	Lepidoptera	Insecta	NA	NL
27	Lemon Pansy	Junonia lemonias	Nymphalidae	Lepidoptera	Insecta	NA	NL
28	Blue Pancy	Junonia orithya	Nymphalidae	Lepidoptera	Insecta	NA	NL

29	Common Bushbrown	Mycalesis perseus	Nymphalidae	Lepidoptera	Insecta	NA	NL
30	Dark Tiger	Tirumala hamata	Nymphalidae	Lepidoptera	Insecta	NA	NL
31	Common Banded Peacock	Papilio crino	Papilionidae	Lepidoptera	Insecta	NA	NL
32	Common Mormon	Papilio polytes	Papilionidae	Lepidoptera	Insecta	NA	NL
33	Mottled Emigrant	Catopsilia pyranthe	Pieridae	Lepidoptera	Insecta	NA	NL
34	Plain Orange Tip	Colotis aurora	Pieridae	Lepidoptera	Insecta	NA	NL
35	Common Jezebel	Delias eucharis	Pieridae	Lepidoptera	Insecta	NA	NL
36	Common Grass Yellow	Eurema hecabe	Pieridae	Lepidoptera	Insecta	NA	NL
37	Yellow Orange Tip	Ixias pyrene	Pieridae	Lepidoptera	Insecta	NA	NL
38	Pale- spotted Emperor	Anax guttatus	Aeshnidae	Odonata	Insecta	LC	NL
39	Golden Dartlet	Ischnura aurora	Coenagrionidae	Odonata	Insecta	NA	NL
40	Indian Common Clubtail	Ictinogomphus rapax	Gomphidae	Odonata	Insecta	LC	NL
41	Common scarlet- darter	Crocothemis erythraea	Libellulidae	Odonata	Insecta	LC	NL
42	Ground Skimmer	Diplacodes trivialis	Libellulidae	Odonata	Insecta	NA	NL
43	Blue Marsh Hawk	Orthetrum glaucum	Libellulidae	Odonata	Insecta	NA	NL
44	Green Marsh Hawk	Orthetrum sabina	Libellulidae	Odonata	Insecta	NA	NL
45	Wandering Glider	Pantala flavescens	Libellulidae	Odonata	Insecta	NA	NL
46	Crimson Marsh Glider	Trithemis aurora	Libellulidae	Odonata	Insecta	NA	NL

Pisces: 14 species of fish belong to 5 orders and 5 families. The order Cypriniformes and Siluriformes each with 5 species in 3 genera of 1 family followed Synbranchiformes (2/1/1), Anabantiformes & Mugiliformes are represented with one species in one genus.

S.No.	Common/Local Name	Scientific Name	Family	Order	IUCN Status	IWPA Schedule
1	Climbing Perch/Goraka	Anabas testudineus	Anabantidae	Anabantiformes	LC	NL
2	Catla	Labeo catla	Cyprinidae	Cypriniformes	LC	NL
3	Rohu/Seelavathi	Labeo rohita	Cyprinidae	Cypriniformes	LC	NL
4	Buradamatta	Channa punctata	Cyprinidae	Cypriniformes	LC	NL
5	Korrameenu	Channa striata	Cyprinidae	Cypriniformes	LC	NL
6	Chittrai	Cirrhinus reba	Cyprinidae	Cypriniformes	LC	NL
7	Corsula	Rhinomugil corsula	Mugilidae	Mugiliformes	LC	NL
8	Long Whiskers Catfish/Jella	Mystus gulio	Bagridae	Siluriformes	LC	NL
9	Errajella	Mystus vittatus	Bagridae	Siluriformes	LC	NL
10	Golden Barb/Perka	Puntius gelius	Bagridae	Siluriformes	LC	NL
11	Buddajella	Rita chrysea	Bagridae	Siluriformes	LC	NL
12	Bondu	Rita kuturnee	Bagridae	Siluriformes	LC	NL
13	Bommidai	Macrognathus guentheri	Mastacembelidae	Synbranchiformes	NA	NL
14	Indian Spiny Eel/Kontemukku	Macrognathus pacalus	Mastacembelidae	Synbranchiformes	LC	NL

Table 16. List of Pisces recorded from the proposed study area

Faunal composition in the study area:

Component	Mammals	Herpetofauna	Birds	Invertebrates	Fishes	Total
Orders	6	3	18	4	5	36
Family	13	13	42	12	5	85
Genera	16	23	70	38	9	156
No. of species	19	26	86	46	14	191

Scheduled species:

- Common Indian Monitor and Indian Peafowl falls under Schedule-I category as per Indian Wildlife Protection Act (IWPA)-1972, Common Indian Monitor also falls under Near Threatened category as per IUCN red list.
- Jungle Cat, Common Fox, Rhesus Macaque, Bonnet Macaque, Hanuman Langur, Indian Chameleon, Checkered Keelback, Indian Cobra, Russell's viper, are the common species that falls under Schedule-II category as per IWPA-1972.

- Nilgai, Indian Spotted Deer, Indian Sambar and Wild Boar falls under Schedule-III category as per IWPA-1972.
- A total of 108 faunal species (Mammals-9, Herpatofauna-15and birds-84) fall under Schedule-IV and Common Crow Schedule-V category according to IWPA- 1972.

Species richness:

Species richness: was high in Avifauna (86 species) followed by Spiders and Butterflies (46 species), Herpetofauna (26 species), Mammals (19 species) and Fishes (14 species), are depicted in Figure 15.



Fig.15. Species richness

The diversity values recorded high for birds (0.108) followed by Invertebrates (0.031), herpetofauna (0.009) Mammals (0.005) and fishes (0.002).

Threatened Fauna Recorded in the Study Area as per IUCN:

Species	IUCN Status
Indian Sambar Rusa unicolor	Vulnerable
Bonnet Macaque Macaca radiata	Vulnerable
Indian Flapshell Turtle Lissemys punctata	Vulnerable
Red sand boa <i>Eryx johnii</i>	Near Threatened
Rough-scaled Sand Boa Gongylophis conicus	Near Threatened
Common Indian Monitor Varanus bengalensis	Near Threatened
Painted Stork Mycteria leucocephala	Near Threatened

Wildlife Conservation Plan:

A Wildlife conservation plan was prepared for conservation of the schedule-I species i.e., Indian Monitor Lizard and Indian Peafowl with an amount of Rs. 478.6 lakhs and was approved by the PCCF (HoFF) & CWLW, Telangana State Forest Department, vide reference No. 5694/2021/Wl-1 dated: 12.10.2021.

Natural streams available in the adjoining forests in 10 Kms radius from the proposed study area:

C N a	No Boat Namo		Streams
5.INO	Beat Name	Nos	Names
1	Tungaram	1	Isuka Vagu
	Sub-Total:	1	
2	Tippanapalli	1	Edumelikala Vagu
	Sub-Total:	1	
3	Penagadapa	1	Tella Vagu
		1	Edumelikala Vagu
	Sub-Total:	2	
4	Ramavaram	1	Tella Vagu
	Sub-Total:	1	
5	Gareebpeta	3	Marri Vagu
			Dargaiah Vagu
			Uppenala Vagu
	Sub-Total:	3	
	Grand Total:	8	

Table 17. List of natural streams in the forest beats providing water to the wildlife inthe 10 Kms radius

Habitat improvement infrastructures & Water Management Initiatives of the Forest Department to sustain and enhance the wildlife population in the 10 Kms radius:

Table 18. No. of fire-line points developed by the Forest Department toavoid/control forest fires within 10 Kms radius

S.No.	Beat Name	Forest Fire Points
1	Tungaram	0
	Sub-Total:	0
2	Tippanapalli	2
	Sub-Total:	2
3	Penagadapa	1
	Sub-Total:	1
4	Ramavaram	0
	Sub-Total:	0
5	Gareebpeta	5
	Sub-Total:	5
	Grand Total:	8

S.No E	Deet News	Percolation Tank (PTs)			
	beat Name	Nos	Latitude	Longitude	
1	Tungaram	1	17.44470	80.62874	
	Sub-Total:	1			
2	Tippanapalli	1	17.43644	80.64424	
		1	17.43769	80.64544	
	Sub-Total:	2			
	Grand Total:	3			

Table 19. No. of PTs developed by the Forest Department

Table 20. List of forest fringe villages within the 10 Kms radius with people depending on the forest for livelihoods and livestock rearing – biotic pressure to existing forest areas:

S No Bost Namo		No of Villages		Develotion	No. of	
5.NO.	Beat Name	Nos.	Names	Population	Cattles	
1	Tungaram	6	Tungaram	3900	2000	
			Sunkara Banjara			
			Tekula Banjara			
			Jarpula Thanda			
			Seemla Thanda			
			Mala Banjara			
	Sub-Total:	6		3900	2000	
2	Tippanapalli	5	Tippanapalli	4400	3200	
			Ahmednagar			
			Venkata Puram			
			Satyanarayana Puram			
			Repallewada			
	Sub-Total:	5		4400	3200	
3	Penagadapa	3	Penagadapa	3000	1500	
			Rampuram			
			Ambedkarnagar			
	Sub-Total:	3		3000	1500	
4	Ramavaram	8	Garimellapadu	4300	280	
			Ramavaram			
			3 Incline			
			2 Incline			
			4 Incline			
			5 Incline			
			Dhanbad			
			Barium Thanda			
	Sub-Total:	8		4300	280	
5	Gareebpeta	15	Gareebpeta	9375	560	
			Laxmipuram Thanda			
			Rudrampur			

		Laxmidevipalli		
		Nimmalagudem		
		Nimmalagudem		
		Colony		
		Marrithanda		
		Chintal Thanda		
		Komatpalli		
		Seethampeta		
		Seethampeta Banjara		
		Reddipalem		
		Ramji Thanda		
		Sujathanagar		
		Mangapeta		
Sub-Total:	15		9375	560
Grand Total:	37		24975	7540

Socio-economic status within the 10 Kms radius from the proposed study area:

Based on the 2011 Census data, in the study area the Schedule caste (SC) population is 16.44%, and the Schedule Tribe (ST) population is 22.08%. Literacy rate in the project area is 66%.

Educational status has improved from generation to generation in this region. Schooling and collages are available up to engineering education. The working population is 41.91% and the non-working population is 58.09%. Infrastructure facilities such as the Road network, Banks, Post office, public distribution system and transport facility and communication facilities are good.

A sample survey of 440 households was surveyed in 15 villages of the study area. The number of females per 1000 males is 1016, showing healthy ratio of female population. Among the surveyed total population, 66% people were literate and 34% were illiterate. This shows that the educational facilities have improved in the sample area.

Distribution of population

As per 2011 census the study area consisted of 2,89,491 persons inhabited in the study area. The distribution of Population in the study area is shown in Table 21.

Particulars	0-2 km	2-5 km	5-10 km	0-10 km
No. of households	4308	4398	66146	74852
Male Population	8623	8394	126544	143561
Female Population	8497	8441	128992	145930
Total Population	17120	16835	255536	289491
Average household size	3.97	3.83	3.86	3.87
Sex ratio	985	1006	1019	1017

Table 21. Distribution of population

*Source: Census 2011

Particulars	0-2 km	2-5 km	5-10 km	0-10 km
Schedule caste	4589	1724	41267	47580
% of SC	26.80	10.24	16.15	16.44
Schedule Tribes	1256	6783	55878	63917
% of ST	7.34	40.29	21.87	22.08
Total SC & ST	5845	8507	97145	111497
% of total SC & ST	34.14	50.53	38.02	38.51
Total Population	17120	16835	255536	289491

Table 22. Distribution of population by social structure

*Source: Census 2011

Household Energy Sources

About 91% of the sample households were using LPG connection, 3% are using both gas and firewood while firewood are used occasionally or in absence of gas. 6% households were using firewood for cooking purpose as they are not having gas connection or immigrants. None of the House Holds are using Kerosene/coal as cooking fuel in the sample households. The LPG connection usage is very high as they are utilising the government subsidy scheme for the LPG connection.

Type of Fuel	No's	%
Gas	401	91
Wood + gas (Dependency on forest)	14	3
Wood (Dependency on forest)	25	6
Total	440	100

Agricultural Scenario

The study area is not agrarian area and there are only a few farmers cultivating Paddy, Mirchi, vegetables and few are growing fodder for their own cattle. 20% of the sample population involved in the other occupation such as services, business, and daily labour works. Sample household survey reveals that only 38% farmers have their own agricultural lands. They were classified as marginal (below 1.00 ha) 37%, small holdings (1-2 ha) 1%, semi-medium no one family found (2 - 4 ha) holdings, medium (4 to 10 ha) and large holdings (10 ha & above) are nil. The remaining 62% doesn't possess any land, as shown in below Table. The major crops cultivated in the study area include Paddy, Mirchi and secondary crop is Maize and millets. The contributions of the agricultural sector to household income are relatively less in 31% of the families are, forming households.

Table	23.	Agriculture	Hodings
-------	-----	-------------	---------

Agriculture land Size Group	Farmers	%
Marginal	162	37
Small	6	1
Small-Medium	0	0
Medium	0	0
Large	0	0
Not having agricultural land	272	62
Total	440	100

Besides land, both in absolute acreage and quality, the ownership of plough bullocks, tractors, and other farm equipment significantly affects the agricultural income. In the study area only 2% possess agricultural related items like tractor and plough. Even though most of them do not possess agricultural related items but they have access to these services through payment/rental basis.

Livestock Farming

Livestock ownership details obtained in the study area during the socio-economic survey for 440 households. A total of 82 households has cattle as secondary source of income. 108 households have Poultry, and 13 families have the Goat/Sheep.

Herd Size	Cow/Buffalo	Goat/Sheep	Poultry
Not having	358	427	332
1 to 3	68	0	86
4 to 7	14	0	19
8 & above	0	13	3
Total	440	440	440

Table 24. Livestock ownership in the 10 Kms around the study area

Infrastructure

Transport facility, mobile communication and postal services are good in the study area. Banks and agricultural societies need to be improved. Women self-help groups are very active and public distribution system is good for ration supply. The details of the facilities are provided below.

Infra		% Cover	age () 201	1		2020 (co	verage %)	
structure Facility	0-2 km	2-5 km	5-10 km	0-10 km	0-2 km	2-5 km	5-10 km	0-10 km
Post Office	50	25	43	39	100	63	74	73
Mobile Phone Coverage	100	100	97	97	100	100	100	100
Public Bus Service	50	75	87	82	100	100	100	100
Black Topped (pucca) Road	100	100	100	100	100	100	100	100
Banks	50	12.5	22	22	100	63	70	70
Self - Help Group	100	100	100	100	100	100	100	100
Public Distribution System	100	100	100	100	100	100	100	100

Table 25. Infrastructure and Educational facilities in the Study area

Table 26. List of floral species documented by the forest department in the adjacentforest areas within the 10 Kms of radius from the proposed study area

Trees			
SI. No	Species	Common Name	
1	Acacia chundra	Sundra	
2	Acacia leucophloea	Tella Thumma	
3	Aegle marmelos	Maredu	
4	Alangium salvifolium	Uduga	
5	Albizia amara	Nalla regu	
6	Albizia lebbeck	Dirishanam	
7	Albizia odoratissima	Chinduga	
8	Andina cordifolia	Bandaru	
9	Anogeissus latifolia	Chirumanu Yelama	
10	Azadirachta indica	Vepa	
11	Bauhinia racemosa	Ari	
12	Bombax ceiba	Buruga	
13	Boswellia serrata	Andugu	
14	Buchanania lanzan	Sarapapu, Mori	
15	Cassia fistula	Rela	
16	Cassine glauca	Bhutangi	
17	Chloroxylon swietenia	Billudu	
18	Cleistanthus collinus	Bankanakkiri, Kodisa	
19	Cochlospermum religiosum	Kondagogu	
20	Cordia dichotoma	Iriki/BankaNakkera	
21	Dalbergia latifolia	Jetregi	
22	Dalbergia paniculata	Soppera/Pachari	
23	Dichrostachys cinerea		
24	Diospyros chloroxylon	Illintha	
25	Diospyros melanoxylon	Tuniki	
26	Erythroxylum monogynum	Devadaru	
27	Eucalyptus tereticornis	Nilagiri	
28	Feronia elephantum	Velaga	
29	Ficus mollis	Juvvi	
30	Ficus racemosa	Medi	
31	Gardenia lucida	Yerra bikki, Karinga	
32	Garuga pinnata	Garga - Garugu	
33	Givotia rottleriformis	Tella poliki	
34	Gmelina arborea	Gummadi Teak	
35	Grewia tiliaefolia	Jana/Thada	
36	Hardwickia binata	Yepa	
37	Holarrhena pubescens	Kodisepala	
38	Holoptelea integrifolia	Nauli	
39	Ixora pavetta	Koravi	
40	Lagerstroemia parviflora	Chennangi	
41	Lannea coromandeliana	Gumpena	
42	Madhuca longifolia	Ірра	

43	Mangifera indica	Mango
44	Manilkara hexandra	Pala/Pedda pala
45	Miliusa tomentosa	Nuluva/ Barre duddi
46	Mitragyna parviflora	Batta ganapa
47	Morinda pubescens	Thogarmogali
48	Naringi crenulata	Torrivelaga
49	Nyctanthes arbor-tristis	Parijatham
50	Paasi	
51	Phyllanthus emblica	Usiri
52	Polyalthia cerasoides	Chilakaduddi
53	Pongamia pinnata	Kanuga
54	Premna tomentosa	Nagur
55	Pterocarpus marsupium	Yegisa
56	Pterospermum xylocarpum	Loluga
57	Schleichera oleosa	Rakot/Pusugu
58	Schrebera swietenioides	Mokkep, Makkam
59	Semecarpus anacardium	Jeedi
60	Soymida febrifuga	Somi
61	Sterculia urens	Thapsi chettu
62	Strychnos nux-vomica	Visha Mushti
63	Strychnos potatorum	Chilla
64	Tamarindus indica	Chintha
65	Tectona grandis	Teak
66	Terminalia alata	Nalla maddi
67	Terminalia arjuna	Tella/Veru maddi
68	Terminalia bellirica	Thani - Thandra
69	Terminalia chebula	Karaka
70	Viscum heyneanum	Karisha
71	Ximenia americana	Udutanakkera
72	Xylia xylocarpa	Bojja/Konda tangedu
73	Ziziphus xylopyrus	Gotti - Gotiki
	Shrubs	
1	Canthium parviflorum	
2	Cissus vitiginea	Gummadi Podha
3	Dodonaea viscosa	Bandaru
4	Grewia hirsuta	Bontha
5	Helicteres isora	Nullthada
6	Jasminum arborescens	Adavi malli
/	Lawsonia inermis	Noillilli
ð	Inaytenus emarginata	Dallulli Sunnomhatti
9	Phyllanthus reticulatus	Summanipatti Manga abattir
10	Kanala spinosa Waadfardig fastisese	Manga chettu
10	woodjorala jruticosa	Jaji Darikathaaga
12	Lizipnus venopiia	rankameega

Climbers				
-	1	Acacia pinnata	Chenchu campa	
	2	Aganasoma caryophyllata	Gudapala	
	3	Asparagus racemosus	Pilli bitiri	
4	4	Cocculus hirsutus	Dusari teega	
ļ	5	Mucuna pruriens	Adavi ulava	
	6	Tylophora indica	Teega Gaddi	
		Herbs		
-	1	Acalypha indica	Muripindi teega	
:	2	Achyranthes aspera	Uttareni/Antisa	
	3	Aerva lanata	Konda pindi	
4	4	Ageratum conyzoides	Goat weed	
ļ	5	Andrographis paniculata	Nelavemu	
	6	Atylosia scarabaeoides	Showy pigeonpea	
,	7 Combretum decandrum		Yedagaddi	
8	8	Curculigo orchioides	Eethakula Gaddi	
	<i>Hemidesmus indicus</i>		Sugandipala	
1	10 Hybanthus enneaspermus		Ratnapurusa	
1	.1	Hyptis suaveolens	Sima tulasi/ Mahaveera	
1	2	Ocimum americanum	Kukka tulasi	
1	3	Ocimum sanctum	Adavi tulasi	
1	4	Phyllanthus niruri	Nela usiri	
1	5	Sida acuta	Chittemu	
		Grasse	S	
1	Aris	tida setacea	Cheepuru gaddi	
2	Cyn	nbopogon martinii	Kasha gaddi	
3	Cynodon dactylon		Garika gaddi	
4	Cyperus pertenius		Nalla Tunga	
5	Сур	erus rotundus	Tunga	
6	6 Dioscorea bulbifera		Adavi gaddu	
7	Ech	inochloa colona	Kaproda gaddi	
8	Ecli	pta alba	Gutta gaddi	
9	Erag	gistielia bifarice		
10	Eragrostis tenella		Chinna Garika Gaddi	

BAMBOOS			
S. No.	Scientific Name	Common Name	
1	Bambusa arundinacea	Mullam Bongu	
2	Dendrocalamus strictus	Sadanam	

Table 26. List of faunal species documented by the forest department in the adjacent forestareas within the 10 Kms of radius from the proposed study area

S.No.	Scientific Name	Common Name
1	Boselaphus tragocamelus	Nilgai
2	Tetracerus qudricornis	Four horned antelope
3	Antilope cervicarpa	Black buck
4	Cervus unicolor	Sambar
5	Axis axis	Spotted deer
6	Sus scrofa	Wild boar
7	Bos gaurus	Indian bison
8	Melurus ursinus	Sloth bear
9	Herpestes edw ardej	Common mongoose
10	Canis aureus	Jackal
11	Vulpes bengalensis	Indian fox
12	Macaca mulatt	Rhesus monkey
13	Presbytis entellus	Common langur
14	Penthera tigris	Tiger
15	Panthera pardus	Leopard
16	Felis chaus	Jungle cat
17	Cuon alpinus	Wild dog

Conclusion

The 313 species of plants and 191 animals that were identified during the brief preliminary survey have highlighted the maximum potential magnitude of biodiversity of the study area. The groups of organisms such as spiders, scorpions, millipedes, insects including butterflies are apparently diverse.

The fauna study reveals the presence of schedule-I faunal species i.e., Indian Monitor lizard, Indian Rock Python and Indian Peafowl falls in the study area of the project. Further, the field survey team have not sighted their presence and did not find any indirect evidence i.e., pug marks, squats, feathers, or dead remains etc. of any wild animals in the study area of the project during field survey. Since, no migratory route / corridors found in the core area. Hence, there is no effect on wildlife migration due to mining.

Recommendations

Restoration and reclamation process during and after mining in the proposed study area and compensatory measures of SCCL with the forest department in the adjoining forests will provide alternate habitats for these species without much competition in resource partition with other existing species around the study area.

These processes should be monitored by a wildlife expert from the beginning of the mining operations, in a long-term period continuously without fail. The long-term monitoring will provide inputs and advisories to maintain and sustain the ecological balance of the study area. One wildlife expert at least for 10 years to monitor the impact of mining in adjoining forest area in the buffer may be engaged.

Also, the monitoring will help in creating a database by conducting the research throughout the mining project covering all seasons as these may also lead to the management of the proposed study area and the adjoining forest areas. Such studies may also raise questions on urban wildlife conservation and its importance in and around the mine operational areas.

A wildlife Conservation/ Mitigation plan was prepared was approved by the Chief Wildlife Warden, TS, vide reference 5694/2021/WL-I Dt.12.10.2021 for Rs.4.78 Crores. This plan shall be implemented within 3 years from the start of mining.

Photo Plates











Butterflies











No.	Common Name	Scientific Name
А.	Tawny Coster	Acraea violae
B.	Chocolate Pansy	Junonia iphita
C.	Common Rose	Pachliopta aristolochiae
D.	Lime	Papilio demoleus
E.	Plain Tiger	Danaus chrysippus
F.	Lemon Pansy	Junonia lemonias
G.	Yellow Orange Tip	Ixias pyrene
H.	Common Grass Yellow	Eurema hecabe
I.	Common Pierrot	Castalius rosimon
J.	Common Indian Crow	Euploea core
К.	Common Banded Awl	Hasora chromu
L.	Indian Cupid	Everes lacturnus
М.	Common Jezebel	Delias eucharis
N.	Common Evening Brown	Melanitis leda

Odonates



No.	Common Name	Scientific Name
А.	Green Marsh Hawk	Orthetrum sabina
В.	Ground Skimmer	Diplacodes trivialis
C.	Golden Dartlet	Ischnura aurora
D.	Ditch Jewel	Brachythemis contaminata
Е.	Common Picture Wing	Rhyothemis variegata
F.	Blue-tailed Green Darner	Anax guttatus

Other Invertebrates



No.	Common Name	Scientific Name
A.	Six-spot Ground Beetle	Anthia sexguttata
B.	Brown Grasshopper	Melanoplus sanguinipes
C.	Painted Grasshopper	Poekilocerus pictus
D.	Palm Rhinoceros Beetle	Oryctes nasicornis
E.	Common Godzilla ant	Componatus compressus
F.	Procession Ant	Leptogenys processionalis

Arachnids



No.	Common Name	Scientific Name
A.	Giant Forest Scorpion	Heterometrus swammerdami
B.	Signature Spider	Argiope anasuja
C.	Common Two Tail	Hersilia savignyi
D.	Decorated Spider	Leucauge decorata
Е.	Yogesh's Green Lynx Spider	Peucetia yogesh
F.	Pawan's Green Lynx Spider	Oxyopes pawani

Reptiles



No.	Common Name	Scientific Name
А.	Indian Garden Lizard	Calotes versicolor
В.	Fan-throated lizard	Sitana ponticeriana
C.	South Indian Rock Agama	Psammophilus dorsalis
D.	Keeled Grass Skink	Eutropis carinata
Е.	Bengal Monitor	Varanus bengalensis
F.	Indian Flap-shelled Turtle	Lissemys punctata
G.	Checkered Keelback	Xenochrophis piscator
H.	Russell's Viper	Daboia russelii

Amphibians



No.	Common Name	Scientific Name
A.	Skittering Frog	Euphlyctis cyanophlyctis
В.	Common Indian Toad	Duttaphrynus melanostictus
C.	Indian Cricket Frog	Fejervarya limnocharis
D.	Indian Bull Frog	Hoplobatrachus tigerinus
Е.	Common Tree Frog	Polypedates maculates
F.	Indian Pond Frog	Euphlyctis hexadactylus
G.	Marbled narrow-mouthed frog	Uperdon variegates
H.	Marbled Balloon Frog	Uperodon systoma

Aves




















No.	Common Name	Scientific Name
A.	Asian Koel	Eudynamys scolopacea
B.	Black Drongo	Dicrurus macrocercus
C.	Red-vented Bulbul	Pycnonotus cafer
D.	Bay-backed Shrike	Lanius vittatus
E.	Spotted Dove	Streptopelia chinensis
F.	Eurasian Golden Oriole	Oriolus oriolus
G.	Cattle Egret	Bubulcus ibis
H.	Green Bee-eater	Merops orientalis
I.	Purple-rumped Sunbird	Nectarinia zeylonica
J.	Brown-capped pygmy woodpecker	Yungipicus nanus
К.	Red-rumped Swallow	Hirundo daurica
L.	Coppersmith Barbet	Psilopogon haemacephalus
М.	White-throated Kingfisher	Halcyon smyrnensis
N.	Common Babbler	Turdoides caudatus
О.	Shikra	Accipiter badius
P .	Scaly-breasted Munia	Lonchura punctulata
Q.	Gray Francolin	Francolinus pondicerianus
R.	Indian Gray Hornbill	Ocyceros birostris
S.	Spotted Owlet	Athene brama
Т.	Pied Bushchat	Saxicola caprata
U.	Rose-ringed Parakeet	Psittacula krameri
V.	Alexandrine Parakeet	Psittacula eupatria
W.	Indian Nightjar	Caprimulgus asiaticus
Χ.	Common Iora	Aegithina tiphia

Mammals



No.	Common Name	Scientific Name
А.	Rhesus Macaque	Macaca mulatta
B.	Bonnet Macaque	Macaca radiata
С.	Hanuman Langur	Semnopithecus entellus
D.	Common Grey Mongoose	Herpestes edwardsii
Е.	Three-striped Palm Squirrel	Funambulus palmarum



Fieldwork Glimpses in the Nature Trails along with the Forest Beat Officers



References

- 1. Sarkar S. Defining Biodiversity. Assessing Biodiversity. The Monist. 2002; 85:131-155.
- 2. Gotelli N, Colwell RK. Quantifying biodiversity: Procedures and pitfalls in the measurement and comparison of species richness. Ecology Letters. 2001; 4:379-391.
- 3. Magurran AE. Meausuring Biological Diversity. Blackwell, 2004.
- 4. Prater SH. The Book of Indian Animals. 3rd Edition. 12th reprint 2005. Bombay Natural History Society, Bombay, 1971, 324.
- 5. Menon VA. Field Guide to Indian Mammals. Dorling Kindersley (India) Pvt. Limited. Delhi, 2003, 1-200.
- 6. Grimmett R, Inskipp C, Inskipp T. Birds of the India, Pakistan, Nepal, Bangladesh, Bhutan, Sri Lanka and the Maldives. Princeton University Press, New Jersey, 2011, 528.
- 7. Ali S. The Book of Indian Birds (13th Revised Edition). Oxford University Press, New Delhi, 2002, 326.
- 8. Smith MA. The Fauna of British India including Ceylon and Burma. Reptilia and Amphibia. Vol. II Sauria. Taylor and Francis, London, 1935, 440.
- 9. Sharma RC. The Reptile fauna of Nagarjunsagar Dam area (Andhra Pradesh, India). Records of Zoological Survey of India. 1971; 63(1-4):77-93.
- 10. Sharma RC. The fauna of India and the adjacent countries Reptilia (Sauria). Zoological Survey of India, Kolkatta, 2002; II:430.sssss
- 11. Whitaker R, Captain A. Snakes of India, The Field Guide. Draco Books, Chennai, 2004; Xiv:481.
- 12. Chanda SK. Handbook: Indian Amphibians. Zoological survey of India, Calcutta, 2002.
- 13. Danial JC. The book of Indian Reptiles and Amphibians. Bombay Natural History Society. 2002; viii+236.
- 14. Ishii M. Transect count of butterflies. In: Decline and conservation of butterflies in Japan 1993; II:91-101.
- 15. Satyarani I. Studies on the Odonates of Andhra Pradesh, India, 9th. Inf. SY111p. odollafol. Madurai 1998; 37-38.
- 16. Fraser, F.C. The Fauna of British India including Ceylon and Burma. Odonata Taylor and Francis Ltd., London, 1933; I:423.
- 17. Fraser FC. The Fauna of British India including Ceylon and Burma. Odonata Taylor and Francis Ltd., London, 1934; II; 398.
- 18. Fraser FC. The Fauna of British India including Ceylon and Burma. Odonata Taylor and Francis Ltd., London, 1936; III:461.
- 19. Kunte K. Butterflies of Peninsular India. Universities Press (Hyderabad) and Indian Academy of Sciences (Bangalore), 2000; xviii+254.
- 20.Issac K. The book of Indian butterflies. Bombay Natural History Society, Bombay. Oxford University Press. 2008, 497.
- 21. Gajbe UA. Fauna of India and the Adjacent Countries: Spiders (Arachnida: Araneae: Oxyopidae). Zoological Survey of India, Kolkata, 2008, 117.
- 22. Tikader BK. Handbook Indian Spiders: A Manual for the study of the Spiders and their relatives-The Scorpions, Pseudoscorpions, Whip scorpions, Harvestmen and all members of the Class Arachnida found in India with analytical keys for their classification and biology. Zoological Survey of India, Kolkata, 1987, 274.
- 23.Sebastian PA, Peter KV. Spiders of India. University Press (India) Private Limited, Hyderabad, 2009.